

# Metformin eligibility up with eGFR versus serum creatinine

November 5 2015

---



(HealthDay)—Use of estimated glomerular filtration rate (eGFR) rather than serum creatinine (sCr) can expand metformin eligibility, according to a study published in the November issue of *Diabetes Care*.

Delphine S. Tuot, M.D., from the University of California in San Francisco, and colleagues examined the potential impact of recommendations to use eGFR rather than sCr to determine metformin eligibility. Metformin eligibility was examined among 3,902 adults with diabetes from the 1999 to 2010 National Health and Nutrition Examination Surveys using conventional sCr thresholds and eGFR categories. Different eGFR equations were used to estimate creatinine clearance (CrCl), including four-variable Modification of Diet in Renal Disease (MDRD), Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) creatinine (CKD-EPIcr), and CKD-EPI cystatin C, as well as

Cockcroft-Gault (CG).

The researchers found that MDRD eGFR  $\geq 45$  mL/min/1.73 m<sup>2</sup> was most common for men and non-Hispanic blacks among adults with sCr above conventional cut-offs (adjusted odds ratios, 33.3 versus women and 14.8 versus whites, respectively). An MDRD eGFR

Citation: Metformin eligibility up with eGFR versus serum creatinine (2015, November 5) retrieved 18 April 2024 from

<https://medicalxpress.com/news/2015-11-metformin-eligibility-egfr-serum-creatinine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.