

Muscle loss linked with falls and fractures in elderly

November 3 2015

Older people with an age-related loss of muscle mass and strength may be at greater risk of falling and bone fractures, according to new research led by the University of Southampton.

A study by an international team of researchers into sarcopenia -where muscles lose form and function with age - found that those with the condition reported higher numbers of falls in the last year and a higher prevalence of fractures.

The decline in muscle mass between the ages of 40 and 80 has been estimated to range from 30 to 60 per cent and is associated with disability, sickness and death. However, there are no universally accepted criteria with which to diagnose patients with sarcopenia and identify those at risk. Three different systems have been suggested by different organisations, as well as a wider term - 'dysmobility syndrome' - which encompasses several adverse factors, including sarcopenia, along with low walking speed and osteoporosis.

"All of these definitions recognise that measuring <u>muscle mass</u> in isolation is inadequate and a measure of muscle function is also required, but there are differences in how function is assessed," says Dr Michael Clynes, lead author of the study from Southampton's Medical Research Council (MRC) Lifecourse Epidemiology Unit. "Due to these differences, an individual may receive different diagnosis depending on how he or she is assessed. This is important because the better we can identify sarcopenia, the more insight we have into the health issues and



risks patients face."

Researchers in the UK, USA and New Zealand assessed a group aged between 70 and 82 using the three sarcopenia definitions, along with the dysmobility syndrome criteria, to see how the different diagnoses corresponded to the occurrence of falls and factures (during the previous year and since the age of 45).

The study, published in Calcified Tissue International, found that the International Working Group on Sarcopenia (IWGS) definition identified the most cases of the condition (8.3 per cent of the cohort) and was linked with significantly higher numbers of falls in the last year and prevalent fractures.

These associations were not apparent when using the European Working Group for Sarcopenia in Older People (EWGSOP) diagnosis or the Foundation for the National Institutes of Health Sarcopenia Project (FNIH) diagnosis (which only identified two per cent as having the condition).

Dysmobility syndrome was common (affecting 24.8 per cent) and corresponded to higher numbers of falls, but no increase in the fracture rate.

"The IWGS definition of sarcopenia appears to be an effective means of identifying individuals at risk of adverse musculoskeletal events, such as falls and fracture," says Dr Clynes. "The findings enable us to more effectively predict those at increased risk of <u>falls</u> and fractures. By defining sarcopenia, health care professionals can target treatment to atrisk individuals."

Professor Cyrus Cooper, Head of MRC Lifecourse Epidemiology Unit, University of Southampton comments: "Such international collaborations



highlight the value of our unique cohort resources in revealing the causes and generating preventive strategies against musculoskeletal ageing."

More information: M. A. Clynes et al. Definitions of Sarcopenia: Associations with Previous Falls and Fracture in a Population Sample, *Calcified Tissue International* (2015). DOI: 10.1007/s00223-015-0044-z

Provided by University of Southampton

Citation: Muscle loss linked with falls and fractures in elderly (2015, November 3) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2015-11-muscle-loss-linked-falls-fractures.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.