

## Noon best time to get vitamin D from sun for minimal cancer risk

November 12 2015



(HealthDay)—Variations in ultraviolet (UV) A and B radiation with increasing latitude and during the day impact skin cancer risk, according to a study published online Nov. 6 in the *International Journal of Dermatology*.

Mantas Grigalavicius, Ph.D., from Oslo University Hospital in Norway, and colleagues examined the latitude dependency of UVB, UVA, vitamin D production, and skin <u>cancer</u> risk in Caucasians. Biologically effective UVB (280 to 315 nm) and UVA (315 to 400 nm) doses were calculated for selected locations. Cancer registries and published articles were reviewed to examine the incidence of <u>squamous cell carcinoma</u> (SCC) and cutaneous melanoma (CM).

The researchers observed a smaller decrease in the annual doses of UVA



radiation versus UVB radiation with increasing latitude. Furthermore, with increasing latitude there were less steep decreases observed in the incidence of CM versus SCC. These data support the role of UVA influencing development of CM. Over one day, the variations in UVA and UVB differed, with the UVB:UVA ratio maximal at noon.

"The best way to obtain a given dose of vitamin D with minimal carcinogenic risk is through a non-burning exposure in the middle of the day, rather than in the afternoon or morning," the authors write.

## More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2015 HealthDay. All rights reserved.

Citation: Noon best time to get vitamin D from sun for minimal cancer risk (2015, November 12) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2015-11-noon-vitamin-d-sun-minimal.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.