

Teen tobacco dependence should be treated with 'same urgency as other drugs,' study says

November 3 2015

Substance abuse treatments that target main issues such as serious drug and alcohol addiction are not frequently being used to also wean adolescents from tobacco, a University of Georgia study finds.

Tobacco addiction in [adolescents](#) is oftentimes an overlooked issue because it doesn't carry with it the stigma that alcohol abuse and other serious drugs do, according to the study's lead author, Jessica Muilenburg, an associate professor at UGA's College of Public Health and health promotion and behavior graduate coordinator.

What most don't realize is that tobacco, she said, "changes the chemistry of your brain and makes you crave whatever your drug of choice is, which is why kicking the tobacco habit with the rest of your addictions is important.

"It's a drug, but it's not treated in the same capacity and with the same urgency as other drugs. We are saying to treat it with the same urgency, because relapse is less likely if you treat the nicotine as well."

Muilenburg has focused much of her research on tobacco use in adolescents and young adults—considering treatment factors that might help them beat these behaviors permanently. For this study, published in the *Journal of Adolescent Health*, she and her co-authors looked at addiction treatments for adolescents and young adults ages 12-28.

Tobacco use is a public health concern that frequently starts during early adolescence and continues across a person's lifespan. According to the Department of Health and Human Services, approximately 2.6 million adolescents smoke cigarettes, and 3.3 million use some sort of tobacco product.

The researchers tracked down [substance abuse](#) centers throughout the U.S. and analyzed treatment practices in those considered adolescent-only clinics. To gather the necessary information, they completed surveys with each of the counselors in those 22 centers.

After analyzing several factors, researchers found that a small number of counselors in these adolescent-only [substance abuse treatment](#) centers actually implemented some sort of tobacco cessation treatment when seeing patients.

Additionally, they found that a majority of these counselors do have the knowledge to implement tobacco cessation treatments and the ability to prescribe medications—such as the patch or nicotine chewing gum—to help adolescents quit. However, counselors don't typically do so for the adolescents they are seeing.

"Their primary goal is getting them off of alcohol and other drugs, but if we can get them off of all drugs, including tobacco, it will be more beneficial for them in the future," Muilenburg said.

Study results show the implementation of these programs needs to be more widely used for [adolescents](#) with [substance abuse](#) issues, she explained.

The researchers also recommend that more counselor training be implemented. Right now, understanding how to treat [tobacco](#)-dependent individuals is not a requirement when counselors obtain their licensure.

Considering [tobacco](#)'s ability to alter the brain and cause relapse, Muilenburg and study co-authors believe it needs to be taken more seriously.

More information: Jessica L. Muilenburg et al. Substance Use Disorder Counselors' Reports of Tobacco Cessation Services Availability, Implementation, and Tobacco-related Knowledge, *Journal of Adolescent Health* (2015). [DOI: 10.1016/j.jadohealth.2015.06.001](https://doi.org/10.1016/j.jadohealth.2015.06.001)

Provided by University of Georgia

Citation: Teen tobacco dependence should be treated with 'same urgency as other drugs,' study says (2015, November 3) retrieved 13 March 2024 from <https://medicalxpress.com/news/2015-11-teen-tobacco-urgency-drugs.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--