

Weekday sleep changes may raise risk of diabetes, heart disease

November 18 2015

Monday mornings could be harmful to your health. Even routine sleep changes such as waking up early for work during the week may raise the risk of developing metabolic problems such as diabetes and heart disease, according to a new study published in the Endocrine Society's *Journal of Clinical Endocrinology & Metabolism*.

Researchers have long recognized that shift work can contribute to metabolic risk because of the continual disruption to the circadian system. Shift workers are more likely to develop metabolic syndrome, coronary [heart disease](#) and Type 2 [diabetes](#) than employees with regular daytime shifts.

Sleep disruption is among the factors that have contributed to rising rates of diabetes and obesity. More than 29 million Americans have diabetes, and 35.1 percent of American adults are obese, according to the Endocrine Society's [Endocrine Facts and Figures report](#).

"Social jetlag refers to the mismatch between an individual's biological circadian rhythm and their socially imposed [sleep](#) schedules. Other researchers have found that social jetlag relates to obesity and some indicators of cardiovascular function," said Patricia M. Wong, MS, of the University of Pittsburgh in Pittsburgh, PA. "However, this is the first study to extend upon that work and show that even among healthy, working adults who experience a less extreme range of mismatches in their sleep schedule, social jetlag can contribute to [metabolic problems](#). These metabolic changes can contribute to the development of obesity,

diabetes and cardiovascular disease."

Researchers examined sleep patterns and cardiometabolic risk in a group of 447 men and women who took part in the Adult Health and Behavior Project Phase 2 study. The participants were between the ages of 30 and 54, and they worked at least 25 hours a week outside the home.

Participants wore a wristband that measured their movement and sleep 24 hours a day for a week. The researchers used questionnaires to assess the participants' diet and exercise habits.

Among the participants, nearly 85 percent had a later halfway point in their sleep cycle - a measurement known as midsleep - on free days compared to work days. The other 15 percent had an earlier midsleep on free days than on work days.

Participants who had a greater misalignment between their sleep schedules on free and work days tended to have poorer cholesterol profiles, higher fasting insulin levels, larger waist circumference, higher body-mass index and were more resistant to insulin than those who had less social jetlag. The association persisted even when the researchers adjusted the measurements to account for variation in other sleep measures and health behaviors such as physical activity and calorie intake.

"If future studies replicate what we found here, then we may need to consider as a society how modern [work](#) and social obligations are affecting our sleep and [health](#)," Wong said. "There could be benefits to clinical interventions focused on circadian disturbances, workplace education to help employees and their families make informed decisions about structuring their schedules, and policies to encourage employers to consider these issues."

More information: "Social Jetlag, Chronotype, and Cardiometabolic

Risk," press.endocrine.org/doi/10.1210/jc.2015-2923

Provided by The Endocrine Society

Citation: Weekday sleep changes may raise risk of diabetes, heart disease (2015, November 18)
retrieved 1 May 2024 from

<https://medicalxpress.com/news/2015-11-weekday-diabetes-heart-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--