

Biochemical clues may predict who develops Alzheimer's disease—and who doesn't

December 21 2015



Diagram of the brain of a person with Alzheimer's Disease. Credit: Wikipedia/public domain.

Investigators have wondered why the brains of some cognitively-intact elderly individuals have abundant pathology on autopsy or significant amyloid deposition on neuroimaging that are characteristic of Alzheimer disease (AD). Researchers reporting in *The American Journal of Pathology* investigated biochemical factors and identified differences in proteins from parietal cortex synapses between patients with and those without manifestation of dementia. Specifically, early-stage AD patients had elevated concentrations of synaptic soluble amyloid- β (A β) oligomers compared to controls who were not demented but displayed signs of AD pathology. Synapse-associated hyperphosphorylated tau (p-



tau) levels did not increase until late-stage AD.

"Our results suggest that effective therapies will need to target synaptic $A\beta$ oligomers, and that anti-amyloid therapies will be much less effective once synaptic p-tau pathology has developed, thus providing a potential explanation for the failure of amyloid-based trials," explained lead investigator Karen H. Gylys, PhD, of the UCLA School of Nursing and the Mary S. Easton Center for Alzheimer's Research at UCLA.

The investigators analyzed brain autopsy samples from different regions in the brain (parietal, superior parietal, entorhinal cortex, and hippocampus) from 46 patients who were classified into groups based on clinical and pathologic criteria: four cognitively-normal elderly controls, two patients with spinocerebellar ataxia type II, 15 patients with no clinical history of dementia but with histopathological signs of ADrelated pathology (high-pathology controls), and 24 patients who were clinically demented and histopathologically diagnosed with AD. Patients with early-stage AD were distinguished from those in later stages. Flow cytometry analysis of synaptosomes (resealed nerve terminals) was used to measure the concentrations of two of the biochemical hallmarks of AD, A β and p-tau, in synaptic terminals.

Investigators examined whether synaptic A β levels were associated with neuritic plaque levels in the <u>parietal cortex</u>. They found little or no evidence of A β immunolabeling in either of the control groups but observed a rise in synaptic A β concentration associated with increasing neuropathologic disease stages. Synaptic A β levels highly correlated with the occurrence of plaque.

Next researchers investigated how $A\beta$ levels are related to clinical dementia. They measured synapse-associated soluble oligomers, known as $oA\beta$ in the parietal cortex, levels of which did not correlate with $A\beta$ plaque counts. However, levels of synaptic $oA\beta$ in early-stage AD, but



not late-stage AD, were significantly elevated relative to both the neuropathologically normal and high pathology groups. "The sharp $oA\beta$ elevation in early AD cases suggests that the clinical syndrome of AD dementia may emerge once the level of synapse-associated soluble oligomers exceeds a certain threshold," noted Dr. Gylys.

Investigators studied the timing of the biochemical changes, noting that other investigators have found evidence that the soluble oligomers of $A\beta$ are the primary toxic peptides that initiate downstream tau pathology as part of the "amyloid cascade hypothesis" of AD. They reported synaptic accumulation of $A\beta$ in the earliest plaque stages, prior to the appearance of synaptic p-tau, which was generally absent until late-stage AD. $A\beta$ and tau levels correlated better in samples from the hippocampus and entorhinal cortex, regions of the brain that are affected earlier in AD than the parietal cortex.

In future work, the authors aim to clarify the precise mechanisms by which soluble A β oligomers affect tau and lead to synaptic dysfunction in AD. An intriguing question is whether therapies that slow A β oligomer accumulation might delay or even prevent the onset of ADrelated dementia. "The correspondence between our human and animal data suggests that this and related animal models will be useful for understanding the progress of synaptic <u>pathology</u> and developing therapies to protect synaptic terminals," commented Dr. Gylys.

More information: "Synaptic Aβ oligomers precede p-tau and differentiate high pathology control cases," by Tina Bilousova, Carol A. Miller, Wayne W. Poon, Harry V. Vinters, Maria Corrada, Claudia Kawas, Eric Y. Hayden, David B. Teplow, Charles Glabe, Ricardo Albay III, Gregory M. Cole, Edmond Teng, Karen Hoppens Gylys (DOI: dx.doi.org/10.1016/j.ajpath.2015.09.018). This article appears online ahead of The *American Journal of Pathology*, Volume 186, Issue 1 (January 2016)



Provided by Elsevier

Citation: Biochemical clues may predict who develops Alzheimer's disease—and who doesn't (2015, December 21) retrieved 8 May 2024 from https://medicalxpress.com/news/2015-12-biochemical-clues-alzheimer-diseaseand-doesnt.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.