

Cell-free circulating mtDNA identifies Parkinson's disease

December 10 2015



(HealthDay)—Cell-free circulating mitochondrial DNA (ccf-mtDNA) from cerebrospinal fluid (CSF) is reduced in patients with Parkinson's disease (PD), according to research published in the December issue of the *Annals of Neurology*.

Angela Pyle, Ph.D., from the University of Newcastle Upon Tyne in the United Kingdom, and colleagues examined whether ccf-mtDNA was an early indicator of PD pathology. Fifty-six CSF samples from patients with PD were selected with 10 age-matched asymptomatic control CSF samples.

The researchers observed a significant reduction in ccf-mtDNA in PD cases versus controls, when analyzing two mtDNA targets: *MTND1* and *MTDN4* copy number. In receiver operating characteristic curve

analysis, *MTND1*- and *MTND4*-calculated ccf-mtDNA strongly predicted PD status (area under curve, 0.81 and 0.84, respectively). There was no significant correlation for ccf-mtDNA with CSF-tau, -phosphorylated tau, and - α -synuclein.

"Given the severity of the reduction in CSF ccf-mtDNA in PD, and supported by both subsequent measurement and remarkably similar data observed in Alzheimer's Disease, we conclude that ccf-mtDNA is a viable biomarker for the early detection of neurodegenerative disease," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Cell-free circulating mtDNA identifies Parkinson's disease (2015, December 10)
retrieved 25 April 2024 from
<https://medicalxpress.com/news/2015-12-cell-free-circulating-mtdna-parkinson-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--