

## **Central venous pressure-guided hydration beneficial in CKD, CHF**

December 11 2015



(HealthDay)—For patients with chronic kidney disease (CKD) and congestive heart failure (CHF) undergoing coronary procedures, central venous pressure (CVP)-guided hydration is associated with reduced risk of contrast-induced nephropathy (CIN), according to a study published online Dec. 9 in *JACC: Cardiovascular Interventions*.

Geng Qian, M.D., from the Chinese People's Liberation Army General Hospital in Peking, and colleagues conducted a prospective comparative trial involving 264 patients with CKD and CHF undergoing coronary procedures. Patients were randomized to CVP-guided hydration or standard hydration (132 patients in each group). The hydration infusion rate was dynamically adjusted according to CVP level every hour in the CVP-guided group.



The researchers found that the total mean volume of isotonic saline administered was significantly higher in the CVP-guided hydration group than the <u>control group</u> (P acute heart failure during the hydration (3.8 versus 3.0 percent; P = 0.500).

"CVP-guided fluid administration can safely and effectively reduce the risk of CIN in patients with CKD and CHF," the authors write.

More information: Full Text (subscription or payment may be required) Editorial (subscription or payment may be required)

Copyright © 2015 <u>HealthDay</u>. All rights reserved.

Citation: Central venous pressure-guided hydration beneficial in CKD, CHF (2015, December 11) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2015-12-central-venous-pressure-guided-hydration-beneficial.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.