

Top risk factors for child undernutrition in India identified

December 17 2015

In India, nearly 40% of all children are stunted—of extremely low height for their age—and nearly 30% are underweight. A new study from Harvard T.H. Chan School of Public Health has now pinpointed the five top risk factors responsible for more than two-thirds of the problem.

The study—the first to comprehensively analyze and estimate the relative importance of known risk factors for child undernutrition—appears online in *Social Science & Medicine*.

Examining an array of 15 well-known risk factors for chronic undernutrition among children in India, the study found that the five top [risk factors](#) were essentially markers of poor socioeconomic conditions as well as poor and insecure nutritional environments in children's households.

Specifically, using data on nearly 29,000 children aged 6-59 months from the 3rd India National Family Health Survey, conducted in 2005-06 (the latest data that is publicly available), the researchers found that the five most important predictors of childhood stunting and underweight were:

- short maternal stature
- a mother with no education
- extreme poverty
- poor dietary diversity
- maternal underweight

Meanwhile, factors such as Vitamin A, breastfeeding, use of iodized salt, improved water and sanitation, and even immunization—all currently high priority interventions in the global discourse on addressing undernutrition—accounted for less than 15% of the cases of undernutrition.

"There is an immediate need to not waste time and resources on short-term and 'doable' interventions," said S V Subramanian, professor of population health and geography and senior author of the study. "While asking people to change behaviors and offering piecemeal solutions might provide some short-term relief, such strategies cannot be substituted for the urgent need to improve food and livelihood security."

More information: Daniel J. Corsi et al. Risk factors for chronic undernutrition among children in India: Estimating relative importance, population attributable risk and fractions, *Social Science & Medicine* (2015). [DOI: 10.1016/j.socscimed.2015.11.014](https://doi.org/10.1016/j.socscimed.2015.11.014)

Provided by Harvard T.H. Chan School of Public Health

Citation: Top risk factors for child undernutrition in India identified (2015, December 17) retrieved 24 March 2023 from <https://medicalxpress.com/news/2015-12-factors-child-undernutrition-india.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.