

## Heart injury reduced after bariatric surgery but not lifestyle intervention

December 9 2015

Heart function in morbidly obese patients returns to normal after bariatric surgery but not after lifestyle intervention, reveals research published today in the *European Journal of Preventive Cardiology*. The results suggest that bariatric surgery2 may reduce the risk of cardiovascular disease in morbidly obese patients.

"Obesity is a global problem that is closely linked to the development of <u>cardiovascular disease</u>," said lead author Dr Magnus N. Lyngbakken, a PhD candidate at the University of Oslo in Norway. "Morbidly <u>obese</u> <u>patients</u> are at high risk of heart attack, heart failure and diabetes."

Approximately one in five western Europeans are obese and have a <u>body</u> mass index (BMI) over 30 kg/m<sup>2</sup>, while around 2 to 3% are morbidly obese (BMI over 35 to 40 kg/m<sup>2</sup>).

The study published today investigated whether weight loss by <u>bariatric</u> <u>surgery</u> could reduce levels of high-sensitivity cardiac troponin I (hs-TnI) in morbidly obese patients, which would indicate reduced cardiac stress and chronic subclinical cardiac injury. Cardiac troponins are traditionally used to diagnose acute ischaemic injury and confirm that a patient had a heart attack. The more recent high-sensitive test also identifies low-grade, chronic cardiac damage in patients who do not have established cardiovascular disease.

Concentrations of hs-TnI were compared between 74 morbidly obese patients who had a type of bariatric <u>surgery</u> called Roux-en-Y gastric



bypass, 62 morbidly obese patients who exercised and restricted their calorie intake, and a control group of 30 people with normal weight.

At the beginning of the study, hs-TnI levels were 2.40 ng/L in the bariatric surgery group, 2.35 ng/L in the lifestyle intervention group, and 0.90 ng/L in the normal weight group. At 12 months after the intervention, hs-TnI levels decreased significantly in the surgery group (p

Citation: Heart injury reduced after bariatric surgery but not lifestyle intervention (2015, December 9) retrieved 27 April 2024 from <u>https://medicalxpress.com/news/2015-12-heart-injury-bariatric-surgery-lifestyle.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.