

Inequalities in alcohol-related mortality in Europe

December 1 2015

Alcohol-related conditions are an important contributing factor to the socioeconomic inequality in total mortality in many European countries, according to a study published this week in *PLOS Medicine*. The database analysis, conducted by Johan Mackenbach at Erasmus University Medical Center, Rotterdam, The Netherlands, and colleagues indicates that in some countries, alcohol-related causes account for 10% or more of the socioeconomic inequality in total mortality among men.

Globally, about three million people die from alcohol-related causes every year. In many European countries, the frequency of drinking and the levels of alcohol consumption are greater in higher socioeconomic groups, whereas binge drinking and other problematic forms of alcohol consumption occur more frequently in lower socioeconomic groups. Mackenbach and [colleagues](#) obtained data on deaths from alcoholic psychosis, dependence, and abuse; alcoholic cardiomyopathy; alcoholic liver cirrhosis; and accidental alcohol poisoning from the mortality registers of 17 European countries. Rates of alcohol-related mortality were higher in individuals with less education or with manual (as opposed to non-manual) occupations in all 17 countries. Absolute inequalities were largest in Eastern Europe, Finland and Denmark. Moreover, the absolute inequalities increased markedly in Hungary, Lithuania, Estonia, Finland, and Denmark because of a rapid rise in alcohol-related mortality in lower socioeconomic groups.

The accuracy of these findings is likely to be affected by the use of routinely collected underlying causes of death. However, these findings

indicate that to reduce socioeconomic inequalities in mortality, health professionals and governments need to introduce interventions and policies designed to counter recent increases in alcohol-related mortality in lower [socioeconomic groups](#). The authors state, "[s]tudies of why such increases [in inequality] have not occurred in countries like France, Switzerland, Spain, and Italy can help in developing evidence-based policies in other European [countries](#)."

More information: Mackenbach JP, Kulhánová I, Bopp M, Borrell C, Deboosere P, Kovács K, et al. (2015) Inequalities in Alcohol-Related Mortality in 17 European Countries: A Retrospective Analysis of Mortality Registers. *PLoS Med* 12(12): e1001909. [DOI: 10.1371/journal.pmed.1001909](#)

Provided by Public Library of Science

Citation: Inequalities in alcohol-related mortality in Europe (2015, December 1) retrieved 27 April 2024 from <https://medicalxpress.com/news/2015-12-inequalities-alcohol-related-mortality-europe.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|