

New study links short sleep to distracted secondary eating and drinking

December 14 2015, by L. Amanda Owens



A recent cross-sectional study conducted by researchers at the University of Alabama at Birmingham explored a new pathway between short sleep—less than seven hours—and obesity.

Gabriel S. Tajeu, DrPH, a postdoctoral fellow in UAB's Department of

Epidemiology, in collaboration with Bisakha Sen, Ph.D., professor in the Department of Health Care Organization and Policy, found "[short sleep](#) is associated with more time spent in secondary eating and, in particular, secondary [drinking](#). This potentially suggests a pathway from short sleep to increased caloric intake in the form of beverages and distracted eating and thus potential increased obesity risk, although more research is needed."

"The association between short sleep and [obesity risk](#) is well-established," Tajeu said. "However, we are looking at whether short sleep is linked to more time spent in secondary eating or drinking, that is, eating or drinking beverages other than water—such as sugar-sweetened beverages—while primarily engaged in another activity, such as television watching."

Using data from 28,150 American adults (55.8 percent female) ranging in ages from 21 to 65 who participated in the American Time Use Survey between 2006 and 2008, the investigators assessed time spent on secondary eating and drinking as well as primary eating and drinking, with [sleep duration](#) as the principal independent variable.

The authors estimated multivariable regression models, an analysis technique that uses multiple variables including participants' demographic characteristics like race and gender, socioeconomic characteristics. and weekday versus weekend participation in ATUS, in order to determine the association of short sleep and eating and drinking behaviors. Compared with participants reporting having experienced normal sleep—between seven and eight hours —those reporting short sleep engaged in secondary eating an additional 8.7 minutes a day as well as an additional 28.6 and 31.28 minutes daily of secondary drinking on weekdays and weekends, respectively.

"New Pathways From Short Sleep to Obesity? Associations Between

Short Sleep and 'Secondary' Eating and Drinking Behavior" was published online in November in the *American Journal of Health Promotion*.

More information: Gabriel S. Tajeu et al. New Pathways From Short Sleep to Obesity? Associations Between Short Sleep and "Secondary" Eating and Drinking Behavior, *American Journal of Health Promotion* (2015). [DOI: 10.4278/ajhp.140509-QUAN-198](https://doi.org/10.4278/ajhp.140509-QUAN-198)

Provided by University of Alabama at Birmingham

Citation: New study links short sleep to distracted secondary eating and drinking (2015, December 14) retrieved 2 May 2024 from <https://medicalxpress.com/news/2015-12-links-short-distracted-secondary.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--