

New method for better treatment of breast cancer

December 30 2015



Mammograms showing a normal breast (left) and a breast with cancer (right).
Credit: Public Domain

A new study shows that a novel imaging-based method for defining appropriateness of breast cancer treatment is as accurate as the current standard-of-care and could reduce the need for invasive tissue sampling. The results suggest that the method might lead to more optimal treatment of individual patients.

Measuring the growth factor HER2 (human epidermal growth factor receptor type 2) is an important tool for deciding correct treatment in [breast cancer](#). Treatments targeting HER2 are expensive but save the lives of many women. Targeted treatments have no effect if the metastases do not express HER2. The current diagnosis of elevated HER2 expression in [metastatic cancer](#) is based on examination of tissue samples obtained by surgery or needle biopsies from the liver, bones and other organs.

The aim of the current study, published in the Open-Access journal *Theranostics*, was to develop a simpler and non-invasive technique, based on whole-body PET/CT imaging, and compare the results of image analysis to the invasive measurements in the same patients. The study included 16 women with on-going treatment of [metastatic breast cancer](#). Twelve had been diagnosed with a HER2-positive primary tumor and four were HER2-negative. All patients were scanned using combined PET/CT and a novel tracer molecule, ABY-025 Affibody, labeled with the short-lived radioactive isotope gallium-68.

The results showed that the amount of HER2-expression in the metastases was accurately measured with the new [method](#). In addition, the amount of HER2-expression in the metastases was frequently found to be different from the [primary tumor](#), leading to a change in therapy in several patients.

The new method might substitute invasive tissue sampling in the near future. Our study resulted in two patients starting therapy and one patient ending therapy with HER2-targeting drugs says Jens Sörensen, PET-researcher and adjunct professor at the Institution of Surgical Sciences, Uppsala University.

The research group now plans a larger study with more participating hospitals to confirm the results with the intention of making the new

method more widely available to patients.

More information: *Theranostics* 2016; 6(2):262-271. [DOI: 10.7150/thno.13502](https://doi.org/10.7150/thno.13502) , www.thno.org/v06p0262

Provided by Uppsala University

Citation: New method for better treatment of breast cancer (2015, December 30) retrieved 24 April 2024 from <https://medicalxpress.com/news/2015-12-method-treatment-breast-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.