

Postpartum family planning services should be a top reproductive health priority

December 11 2015

Approximately one-quarter of inter-birth intervals in low- and middle-income countries are less than 24 months in length, exposing infants to risks of prematurity, low birthweight, and death. Increased evidence of these health risks has emerged in the past few decades and, after a period of neglect, interest in postpartum family planning has followed, supported by organizations like WHO and USAID.

How successful have interventions and projects to improve postpartum [contraceptive](#) uptake been? A new study published in *Studies in Family Planning* as part of the December special issue "Postpartum and Post-Abortion Contraception: From Research to Programs" analyzes data from 35 studies (19 of which were not included in earlier reviews) to address this important question, and identifies program implications and future research priorities.

Key findings include that counseling before discharge from a maternity unit likely has an impact on subsequent contraceptive use and that the integration of [family planning](#) into immunization and pediatric services is justified (despite few programs offering these services).

The case can also be made for relaxing the strict conditions for LAM (the Lactational Amenorrhea Method based on the premise that breastfeeding interferes with the release of the hormones needed to trigger ovulation) to be considered an effective postpartum contraceptive method.

The authors note: "In view of the findings, the ideal strategy is to incorporate contraceptive advice and services across the continuum of reproductive healthcare." Four types of interventions are analyzed—antenatal, postnatal, combined ante- and postnatal, and integration with other services—and a range of countries in Africa, Asia, and Latin America are included.

More information: John Cleland et al. Interventions to Improve Postpartum Family Planning in Low- and Middle-Income Countries: Program Implications and Research Priorities, *Studies in Family Planning* (2015). [DOI: 10.1111/j.1728-4465.2015.00041.x](https://doi.org/10.1111/j.1728-4465.2015.00041.x)

Provided by Population and Development Review

Citation: Postpartum family planning services should be a top reproductive health priority (2015, December 11) retrieved 5 May 2024 from <https://medicalxpress.com/news/2015-12-postpartum-family-reproductive-health-priority.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
