

Significant lowering of alar rim with cartilage repositioning

December 3 2015



(HealthDay)—Patients who undergo lateral crural repositioning have significant lowering of the alar rim, according to a study published online Nov. 19 in *JAMA Facial Plastic Surgery*.

Christian A. Paquet, M.D., from the University of Southern California in Los Angeles, and colleagues retrospectively reviewed data from 54 [patients](#) (102 hemi-noses; 79 percent women) who had lateral crural [repositioning](#) performed by the same surgeon. Patients were included if they had standardized photographs taken preoperatively and at least six months postoperatively (average time, 11.3 months).

Eighty-three percent of the cases were revision rhinoplasties. The researchers found that there was a net decrease of 6.8 degrees based on the preoperative and postoperative mean anterior nostril apex angles (P

degrees, respectively, were seen when examining patients who underwent lateral crural repositioning alone and those who also had lateral crural strut grafts (both P

"The results demonstrate that repositioning of the lower lateral cartilages with or without a lateral crural strut graft into a more caudal pocket results in a statistically significant lowering of the alar rim," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Significant lowering of alar rim with cartilage repositioning (2015, December 3)
retrieved 2 May 2024 from
<https://medicalxpress.com/news/2015-12-significant-lowering-alar-rim-cartilage.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
