


The ability to control dreams may help us unravel the mystery of consciousness

January 7 2016, by Dan Denis And Giulia Poerio, University Of Sheffield



Credit: Alexey Komissarov from Pexels



We spend around <u>six years of our lives dreaming</u> – that's 2,190 days or 52,560 hours. Although we can be aware of the perceptions and emotions we experience in our dreams, we are not conscious in the same way as when we're awake. This explains why we can't recognise that we're in a dream and often mistake these bizarre narratives for reality.

But some people – lucid dreamers – have the ability to experience awareness during their dreams by "re-awakening" some aspects of their waking <u>consciousness</u>. They can even take control and act with intention in the dream world (think Leonardo DiCaprio in the film Inception).

Lucid dreaming is still an understudied subject, but recent advances suggest it's a hybrid state of waking consciousness and sleep.

Lucid dreaming is one of many "anomalous" experiences that can occur during sleep. Sleep paralysis, where you wake up terrified and paralysed while remaining in a state of sleep, is another. There are also <u>false</u> <u>awakenings</u>, where you believe you have woken up only to discover that you are in fact dreaming. Along with lucid dreams, all these experiences reflect an increase in subjective awareness while remaining in a state of sleep. To find out more about the transitions between these states – and hopefully consciousness itself – we have launched a <u>large-scale online</u> <u>survey</u> on sleep experiences to look at the relationships between these different states of hybrid consciousness.

Lucid dreaming and the brain

About <u>half of us</u> will experience at least one lucid dream in our lives. And it could be something to look forward to because it allows people to simulate desired scenarios from meeting the love of their life to winning a medieval battle. There is some evidence that <u>lucid dreaming can be induced</u>, and a number of large online communities now exist where users share tips and tricks for achieving greater lucidity during their



dreams (such as having dream totems, a familiar object from the waking world that can help determine if you are in a dream, or spinning around in dreams to stop lucidity from slipping away).



Sleep paralysis. My Dream, My Bad Dream, 1915. Credit: Fritz Schwimbeck/wikimedia

A <u>recent study</u> that asked participants to report in detail on their most recent dream found that lucid (compared to non-lucid) dreams were indeed characterised by far greater insight into the fact that the sleeper was in a dream. Participants who experienced lucid dreams also said they had greater control over thoughts and actions within the dream, had




the ability to think logically, and were even better at accessing real memories of their waking life.

Another study looking at people's ability to make conscious decisions in waking life as well as during lucid and non-lucid dreams found a large degree of overlap between volitional abilities when we are awake and when we are having lucid dreams. However, the ability to plan was considerably worse in lucid dreams compared to wakefulness.

Lucid and non-lucid dreams certainly feel subjectively different and this might suggest that they are associated with different patterns of brain activity. But confirming this is not as easy as it might seem. Participants have to be in a brain scanner overnight and researchers have to decipher when a lucid dream is happening so that they can compare brain activity during the lucid dream with that of non-lucid dreaming.

Ingenious studies examining this have devised a communication code between lucid dreamer participants and researchers during Rapid Eye Movement (REM) sleep, when <u>dreaming typically takes place</u>. Before going to sleep, the participant and the researcher agree on a specific eye movement (for example two movements left then two movements right) that participants make to signal that they are lucid.





Prefrontal cortex. Credit: Natalie M. Zahr, Ph.D., and Edith V. Sullivan, Ph.D. - Natalie M. Zahr, Ph.D., and Edith V. Sullivan, Ph.D.

By using this approach, studies have found that the shift from non-lucid to lucid REM sleep is associated with an increased activity of the frontal areas of the brain. Significantly, these areas are associated with "higher order" cognitive functioning such as logical reasoning and voluntary behaviour which are typically only observed during waking states. The type of brain activity observed, gamma wave activity, is also known to allow different aspects of our experience; perceptions, emotions, thoughts, and memories to "bind" together into an integrated consciousness. A follow-up study found that electrically stimulating these areas caused an increase in the degree of lucidity experienced during a dream.



Another study more <u>accurately specified the brain regions</u> involved in lucid dreams, and found increased activity in regions such as the prefrontal cortex and the precuneus. These brain areas are associated with higher cognitive abilities such as <u>self-referential processing</u> and a sense of agency – again supporting the view that lucid dreaming is a hybrid state of consciousness.

Tackling the consciousness problem

How consciousness arises in the brain is one of the most perplexing questions in neuroscience. But it <u>has been suggested</u> that studying <u>lucid dreams</u> could pave the way for new insights into the neuroscience of consciousness.

This is because lucid and non-lucid REM sleep are two states where our conscious experience is markedly different, yet the overall brain state remains the same (we are in REM sleep all the time, often dreaming). By comparing specific differences in brain activity from a lucid dream with a non-lucid one, then, we can look at features that may be facilitating the enhanced awareness experienced in the lucid dream.

Furthermore, by using eye signalling as a marker of when a sleeper is in a lucid dream, it is possible to study the neurobiological activity at this point to further understand not only what characterises and maintains this heightened consciousness, but how it emerges in the first place.

This article was originally published on The Conversation. Read the original article.

Source: The Conversation



Citation: The ability to control dreams may help us unravel the mystery of consciousness (2016, January 7) retrieved 4 May 2024 from https://medicalxpress.com/news/2016-01-ability-unravel-mystery-consciousness.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.