

Sleep apnea treatment associated with reduced readmissions for patients with heart failure

January 26 2016




Illustration of obstruction of ventilation. Credit: Habib M'henni / public domain

Early diagnosis and treatment of sleep apnea may reduce six-month readmissions for patients hospitalized with heart failure, [according to research recently published online](#) by the *American Journal of Cardiology*.

"Our research showed that early recognition and treatment of patients hospitalized with decompensated congestive [heart failure](#) is associated with a reduction in readmissions, for patients who use their positive airway pressure (PAP) therapy on a regular basis," said first author Sunil Sharma, M.D., FAASM, Associate Professor of Pulmonary Medicine in the Sidney Kimmel Medical College at Thomas Jefferson University. "Importantly, hospitals can implement cost-effective screening programs to catch [sleep apnea](#) in hospitalized, high risk patients."

Dr. Sharma and the team screened patients admitted to the hospital with heart failure, for underlying sleep disordered breathing. Of the 75 patients that followed up with an outpatient polysomnography, the gold standard test, 70 received the diagnosis of sleep disordered breathing. Over the next six months, the team tracked patients' PAP compliance, emergency room visits and readmissions. Compliance was monitored objectively by the device computer and defined as a minimum of four hours of use, 70 percent of the time, for four weeks consecutively or more during the first three months of therapy.

By comparing pre- and post-treatment readmissions in compliant (n=37) and non-compliant patients (n=33), the researchers found a reduction in hospital visits for those who used their PAP regularly over a period of six months.

"Physicians should be on the lookout for sleep apnea in [patients](#) with heart failure with the goal of diagnosing and treating early, which might help prevent readmissions and [emergency room visits](#)," Sharma said. "Multi-center, randomized studies are needed to replicate and validate

these findings."

More information: Sunil Sharma et al. Effect of Early Intervention with Positive Airway Pressure Therapy for Sleep Disordered Breathing on Six-Month Readmission Rates in Hospitalized Patients with Heart Failure, *The American Journal of Cardiology* (2015). [DOI: 10.1016/j.amjcard.2015.12.032](https://doi.org/10.1016/j.amjcard.2015.12.032)

Provided by Thomas Jefferson University

Citation: Sleep apnea treatment associated with reduced readmissions for patients with heart failure (2016, January 26) retrieved 25 April 2024 from <https://medicalxpress.com/news/2016-01-apnea-treatment-readmissions-patients-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.