

# Attention neuron type identified

January 14 2016

---



Marie Carlén, Ph.D., is a principal investigator at the Department of Neuroscience, Karolinska Institutet in Sweden. Credit: Gustav Mårtensson

Researchers at Karolinska Institutet in Stockholm, Sweden, have identified for the first time a cell type in the brain of mice that is integral to attention. By manipulating the activity of this cell type, the scientists were able to enhance attention in mice. The results, which are published

in the journal '*Cell*', add to the understanding of how the brain's frontal lobes work and control behaviour.

The frontal cortex of the brain plays a crucial part in cognitive functions, including everyday mental processes such as attention, memory, learning, decision-making and problem-solving. However, little is known about how the frontal cortex performs these [mental processes](#), including which neuronal cell types are involved. A longstanding theory holds that parvalbumin-expressing neurons (PV cells) play a key role in cognition; now, however, researchers show that PV cells seem to be not only essential to attention; it also appears that it is enough to optimise PV cell [activity](#) in order to enhance attention.

The team focused on attention since it is a cognitive process that is affected in many neuropsychiatric disorders. The scientists trained mice to perform a task requiring a high degree of attention, and recorded the activity of hundreds of individual neurons in the [frontal cortex](#) while the animals repetitively performed the task.

"We found that the activity of the PV cells reflected the animals' level of attention," says lead investigator Marie Carlén at the Department of Neuroscience. "The PV cells were highly active if the animals were attentive, and less active when they were inattentive. The differences were so great that we were able to predict if an animal would perform the task successfully or not merely by looking at the activity of the PV cells."

The researchers used optogenetics to influence PV-cell activity during the seconds the animals needed to be attentive. They found that the animals' attention was impaired when they either inhibited or changed the pattern of this activity. However, they also found a type of manipulation that could improve [attention](#). During certain cognitive processes a category of brain waves known as gamma oscillations (30-80

Hz) increases in prefrontal cortex, and when the scientists activated the PV cells at gamma frequencies the animals solved the task more times.

While cognitive problems are common in mental disorders such as schizophrenia, ADHD and autism, there is currently no effective medicine available.

"Our findings tie together a number of previous observations on PV cells and their involvement in cognition and [neuropsychiatric disorders](#), and demonstrate this cell type's critical role in cognition and psychiatry," says Dr Carlén. "The findings also show that it's possible to enhance cognitive functions by altering the activity of a single neuron type, which is quite astonishing when you think of how complex the brain is. PV [cells](#) are therefore a very interesting target for the pharmaceutical industry."

**More information:** 'Prefrontal Parvalbumin Neurons in Control of Attention', Hoseok Kim, Sofie Åhrlund-Richter, Xinming Wang, Karl Deisseroth, Marie Carlén, *Cell*, online 14 January 2016, doi: [dx.doi.org/10.1016/j.cell.2015.11.038](https://doi.org/10.1016/j.cell.2015.11.038)

Provided by Karolinska Institutet

Citation: Attention neuron type identified (2016, January 14) retrieved 5 May 2024 from <https://medicalxpress.com/news/2016-01-attention-neuron.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---