

Frailty may increase complication risk following urologic surgery

January 19 2016

For patients undergoing urologic surgery, frailty may increase their risk of experiencing complications after surgery.

Among 95,108 patients undergoing 21 different urologic procedures, such as bladder or [prostate removal](#), the average frequency of [complications](#) per individual was 11.7%, with the most common complications being hospital readmission (6.2%), blood transfusion (4.6%), and urinary tract infection (3.1%).

Tests of frailty revealed that patients who were frail, a physical vulnerability that can lead to increased susceptibility to disability, were 1.7-times more likely to experience complications after surgery.

"We found that frailty was a significant predictor of postoperative complications independent of age and across most procedures evaluated, making it a very important factor to consider in preoperative decision-making and counseling," said Dr. Anne Suskind, lead author of the *BJU International* study.

More information: Anne M Suskind et al. The impact of Frailty on complications in patients undergoing common urologic procedures; a study from the American College of Surgeons National Surgical Quality Improvement Database, *BJU International* (2015). [DOI: 10.1111/bju.13399](#)

Provided by Wiley

Citation: Frailty may increase complication risk following urologic surgery (2016, January 19) retrieved 20 March 2024 from <https://medicalxpress.com/news/2016-01-frailty-complication-urologic-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.