

Exercise for people with dementia improves balance and reduces dependence

January 20 2016







Regular exercise improves balance for people with dementia and reduces dependence on assistance. Credit: MostPhotos

Regular exercise improves balance for people with dementia and reduces dependence on assistance. This according to new research on healthcare for people suffering from dementia conducted at Umeå University in Sweden, which has now been published in the *Journal of the American Geriatrics Society*.

In a unique study on people with dementia living in residential care facilities, researchers from Umeå University have found that regular functional exercise, similar to everyday activities and performed at high intensity, can improve balance and reduce dependence on assistance in activities of daily living, such as for instance mobility or toilet visits. Training sessions lasting 45 minutes, two to three times per week, can lead to an improved quality of life for individuals suffering from dementia – a progressive illness leading to gradual reduction in cognitive and physical function.

In 2012, the number of individuals with dementia in Sweden stretched to 150,000 according to the National Board of Health and Welfare (Socialstyrelsen). This is a number expected to reach at least 250,000 in 2050 at the speed of population ageing. Having an increased proportion of individuals with dementia in society increases the demand and costs for healthcare. At present, the cost of dementia in Sweden reaches around SEK 60 billion per annum, according to the National Board of Health and Welfare.

"Regular exercise has a positive effect on people with dementia and



should, therefore, be included in the care in residential care facilities. Studies such as the present one are rare, but provide important knowledge to further build upon in order to develop care of people with dementia as a cost-effective means of meeting future challenges, and help individuals to maintain independence longer," says Annika Toots, PhD student at the Department of Community Medicine and Rehabilitation at Umeå University, and first author of the article.

The study, described in the article Effects of a high-intensity functional exercise program on dependence in activities of <u>daily living</u> and balance in older adults with dementia, is a part of the greater Umeå Dementia and Exercise Study (UMDEX), performed involving people with dementia in <u>residential care</u> facilities. As a next step, research groups at the Unit of Physiotherapy and Geriatric medicine will analyse which effect exercise has on for instance ability to walk and cognitive ability in the target group, as well as how participants and physiotherapists experienced the exercise programme.

The published study involved 186 people with dementia in 16 different residential <u>care facilities</u> in the Umeå area. All participants were 65 years of age or above and in need of personal care. The participants were randomly allocated into two groups, where one group undertook a high-intensive, functional exercise programme led by physiotherapists. The programme included various functional exercises that aimed to improve leg strength, balance and walking, which are part of everyday activities. The workout covered 45 minutes, two to three times per week for a duration of four months.

Instead of training, the sedentary control group took part in stimulating activities of group conversations, singing and reading aloud sessions. The purpose of this was to control the positive effects that stimulation through togetherness and attention have. All participants were tested before, as well as four and seven months after the completion of the



programme.

Due to the progressive course of dementia, a deterioration was noted in all participants' abilities to independently manage <u>everyday activities</u>. The deterioration occurred at a slower pace in the exercise group and they showed an improved balance. The positive effects of the exercise varied depending on the type of dementia, where the group with <u>vascular dementia</u> experienced better effects of exercising than participants with Alzheimer's disease. To better plan and carry out <u>exercise</u> for <u>people</u> with dementia, it can, therefore, be of importance to identify the type of <u>dementia</u>.

More information: Annika Toots et al. Effects of a High-Intensity Functional Exercise Program on Dependence in Activities of Daily Living and Balance in Older Adults with Dementia, *Journal of the American Geriatrics Society* (2016). DOI: 10.1111/jgs.13880

Provided by Umea University

Citation: Exercise for people with dementia improves balance and reduces dependence (2016, January 20) retrieved 26 April 2024 from <u>https://medicalxpress.com/news/2016-01-people-dementia.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.