


Twin study estimates familial risks of 23 different cancers

January 5 2016



A large new study of twins has found that having a twin sibling diagnosed with cancer poses an excess risk for the other twin to develop any form of cancer. Among the 23 different types of cancer studied, an excess familial risk was seen for almost all of the cancers, including common cancers such as breast and prostate cancer, but also more rare

cancers such as testicular cancer, head and neck cancer, melanoma, ovarian and stomach cancer.

The study, led by researchers at the Harvard T.H. Chan School of Public Health, the University of Southern Denmark, and the University of Helsinki, is the first to provide family risk estimates for these and other rarer cancers. The study also showed, for the first time, that in twin pairs where both developed cancer, each twin often developed a different type of cancer—which suggests that, in some families, there is a shared increased risk of any type of cancer.

"Prior studies had provided [familial risk](#) and heritability estimates for the common cancers—breast, prostate, and colon—but, for rarer cancers, the studies were too small, or the follow-up time too short, to be able to pinpoint either heritability or family risk," said Lorelei Mucci, associate professor of epidemiology at Harvard Chan School and co-lead author of the study.

The study will be published online January 5, 2016 in *JAMA (Journal of the American Medical Association)*.

Familial risk of cancer is a measure of the [cancer risk](#) in an individual. The study also looked at heritability of cancer, a measure of how much of the variation in cancer risk of populations is due to genetic factors.

"Findings from this prospective study may be helpful in patient education and cancer risk counseling," said Jaakko Kaprio, from the University of Helsinki and co-senior author of the study.

The researchers looked at more than 200,000 [twins](#), both identical and fraternal, in Denmark, Finland, Norway, and Sweden, who participated in the Nordic Twin Study of Cancer followed over an average of 32 years between 1943 and 2010. Large twin studies can help scientists

assess the relative contribution of inherited factors in cancer and characterize familial cancer risk by taking into account the genetic relatedness of identical and fraternal twins.

Overall, one in three people in the study developed cancer over the course of a lifetime. Cancer was diagnosed in both twins for 3,316 of the pairs, in whom the same cancer was diagnosed among 38% of the identical twins and 26% of the fraternal twins. The researchers estimated that, when one fraternal twin was diagnosed with any cancer, the co-twin's risk of getting cancer was 37%; among identical twins, the risk jumped to 46%. One of the strongest familial risks was observed for testicular cancer. The researchers found that a man's risk of developing this disease was 12 times higher if his fraternal twin developed it, and 28 times higher if his genetically identical twin developed it.

Given the fact that fraternal twins are similar genetically to siblings who aren't twins, the finding of excess cancer risk among fraternal twin pairs can provide information about an increased cancer risk for families in which one sibling gets cancer.

The researchers also found that the heritability of cancer overall was 33%. Significant heritability was found for skin melanoma (58%), prostate cancer (57%), non-melanoma skin cancer (43%), ovarian cancer (39%), kidney cancer (38%), breast cancer (31%), and uterine cancer (27%).

"Because of this study's size and long follow-up, we can now see key genetic effects for many cancers," said Jacob Hjelmberg, from the University of Southern Denmark and co-lead author of the study.

"This study was possible given the unique databases in the four Nordic countries, and will be a future resource to solve other complex questions in [cancer](#)," said Hans-Olov Adami, adjunct professor of epidemiology at

Harvard Chan School and co-senior author of the study.

More information: "Familial risk and heritability of cancer among twins in Nordic countries," Lorelei A. Mucci, Jacob B. Hjelmborg, Jennifer R. Harris, Kamila Czene, David J. Havelick, Thomas Scheike, Rebecca E. Graff, Klaus Holst, Sören Möller, Robert H. Unger, Christina McIntosh, Elizabeth Nuttall, Ingunn Brandt, Kathryn L. Penney, Mikael Hartman, Peter Kraft, Giovanni Parmigiani, Kaare Christensen, Markku Koskenvuo, Niels V. Holm, Kauko Heikkilä, Eero Pukkala, Axel Skytthe, Hans-Olov Adami, Jaakko Kaprio, on behalf of the Nordic Twin Study of Cancer (NorTwinCan) collaboration. *JAMA*, online January 5, 2016, [DOI: 10.1001/jama.2015.17703](https://doi.org/10.1001/jama.2015.17703).

Provided by Harvard T.H. Chan School of Public Health

Citation: Twin study estimates familial risks of 23 different cancers (2016, January 5) retrieved 9 April 2024 from <https://medicalxpress.com/news/2016-01-twin-familial-cancers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
