


# Diabetes drug may prevent recurring strokes

February 17 2016

---



Ischemic stroke occurs when a brain blood vessel gets blocked. The gray area represents brain tissue that is not receiving nutrients as a result of the stroke. Credit: NINDS.

Pioglitazone, a drug used for type 2 diabetes, may prevent recurrent stroke and heart attacks in people with insulin resistance but without diabetes. The results of the Insulin Resistance Intervention after Stroke (IRIS) trial, presented at the International Stroke Conference 2016 in Los Angeles and published in the *New England Journal of Medicine*, suggest a potential new method to prevent stroke and heart attack in high-risk patients who have already had one stroke or transient ischemic

attack. This large, international study was supported by the National Institutes of Health's National Institute of Neurological Disorders and Stroke (NINDS).

The IRIS trial is the first study to provide evidence that a drug targeting cell metabolism may prevent secondary strokes and heart attacks even before diabetes develops. Insulin regulates metabolism and keeps blood sugar levels from getting too high, along with many other processes, in the body. Insulin resistance is a condition in which the body produces insulin but does not use it effectively.

"This study represents a novel approach to prevent recurrent vascular events by reversing a specific metabolic abnormality thought to increase the risk for future heart attack or stroke," said Walter J. Koroshetz, M.D., director of the NINDS.

"The IRIS trial supports the value of more research to test the vascular benefits of other interventions such as exercise, diet and medications that have similar effects on metabolism as [pioglitazone](#)," said Walter N. Kernan, M.D. professor of general medicine at Yale University School of Medicine, New Haven, Connecticut, and lead author of the study.

More than 3000 patients from seven countries who had experienced an [ischemic stroke](#) or transient ischemic attack within the previous six months were randomized to receive pioglitazone or placebo for up to five years in addition to standard care. Ischemic stroke and [transient ischemic attacks](#) can occur when a cerebral blood vessel becomes blocked, cutting off the delivery of oxygen and nutrients to brain tissue.

In this study, stroke or heart attack occurred in 9 percent of participants taking pioglitazone and 11.8 percent of patients on placebo, which was a relative decrease of 24 percent. The results suggest that 28 strokes or heart attacks may be prevented for every 1000 patients who take

pioglitazone for up to five years.

Insulin resistance is a hallmark of type 2 diabetes but also occurs in more than 50 percent of people with ischemic stroke who do not have diabetes. People with diabetes are known to have increased risk of stroke. Previous research suggested that [insulin resistance](#) increases risk for stroke, but the IRIS trial was the first to treat it and suggested that the therapy reduced the risk of [recurrent stroke](#) and heart attacks. However, pioglitazone is not FDA-approved for the uses studied in the IRIS trial.

In this study, pioglitazone also reduced the risk of diabetes by 52 percent in the study participants.

The study evidenced an additional known side effect of the drug, which is an increased risk of bone fractures. To help doctors and patients choose the best strategy for preventing recurring strokes, future studies will attempt to identify a person's risk of bone fractures due to pioglitazone. As approved for use in medical practice, the drug also carries additional side effects (drug label).

"More research is needed to determine the mechanisms by which pioglitazone decreases risk for [stroke](#) and [heart attack](#) and increases bone fracture risk, with the hope of developing strategies that maximize benefit and minimize serious side effects in our patients," said Dr. Kernan.

**More information:** Walter N. Kernan et al. Pioglitazone after Ischemic Stroke or Transient Ischemic Attack, *New England Journal of Medicine* (2016). [DOI: 10.1056/NEJMoa1506930](https://doi.org/10.1056/NEJMoa1506930)

Provided by National Institutes of Health

Citation: Diabetes drug may prevent recurring strokes (2016, February 17) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-02-diabetes-drug-recurring.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.