

Research links prenatal stress to babies' health in war zones

February 2 2016

Children from war-torn areas of the globe are affected by trauma even before they are born, according to a new University of Florida study.

To gather their results, researchers went to the eastern Democratic Republic of Congo, a region routinely called "the worst place in the world" to be a woman, said Darlene A. Kertes, the study's lead author and an assistant professor in UF's department of psychology. Women in this unstable region are routinely the target of rape and other war-related traumas.

"Our research shows that stressful life experiences affect our bodies all the way down to our genes," said Kertes, who also is affiliated with the University of Florida Genetics Institute.

The study was published in the January/February issue of *Child Development*, the flagship journal for the Society for Research in Child Development. The results showed that mothers' stressful life experiences were linked with epigenetic markers in key genes that regulate the body's response to <u>stress</u>, in both mothers and newborns.

"The study is one of the first of its kind to be conducted in a developing country," Kertes said. "Most information to date about effects of stress and trauma on prenatal development has been gathered in a Western context."

Samples of <u>umbilical cord blood</u>, placenta and the mothers' blood were



collected at birth and tested for impacts of war trauma and <u>chronic stress</u>. The researchers looked at DNA methylation, an epigenetic process that makes genes more or less able to respond to biochemical signals in the body.

During pregnancy, a mother's bodily responses to stress are passed onto the fetus, affecting a child's brain development, birth weight and functioning of the children's own HPA axis even after they are born.

The researchers looked at the babies' birth weight as an indicator of children's overall development. They found that stress-linked DNA methylation differences predicted lower <u>birth weight</u>.

"The <u>stress exposure</u> affected the maternal and fetal tissues differently, which shows that the impact of stress differs depending on an individual's life phase," Kertes said, adding that stress experienced at very young ages affects the way the body responds to stress throughout life.

This is the first time researchers have documented stress effects, either pre- or postnatal, on methylation of a gene called CRH in humans. CRH makes a hormone that triggers the body's stress response. The study also confirmed stress effects on several other genes known to be involved in the <u>stress response</u>.

Kertes and her colleagues have started to examine the longer term effects of stress on <u>child development</u> in conflict-ridden regions. She emphasized that traumatic events can also have cross-generational impacts.

"War and conflict do not just impact the health and well-being for people who experience it directly," she said. "It can potentially have longterm consequences for future generations."



Provided by University of Florida

Citation: Research links prenatal stress to babies' health in war zones (2016, February 2) retrieved 6 May 2024 from https://medicalxpress.com/news/2016-02-links-prenatal-stress-babies-health.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.