


# **Omega-3 fatty acids may lower breast cancer risk in postmenopausal obese women**

February 24 2016

---



Micrograph showing a lymph node invaded by ductal breast carcinoma, with extension of the tumour beyond the lymph node. Credit: Nephron/Wikipedia

Omega-3 fatty acids may lower the risk of breast cancer in postmenopausal obese women, according to researchers.

The protection likely comes from the [fatty acids](#)' anti-inflammatory effects, said Andrea Manni, professor and division chief of endocrinology, diabetes and metabolism, Penn State College of Medicine.

Obesity is a major [breast cancer](#) risk factor in [postmenopausal women](#), and scientists believe increased inflammation is an important underlying cause in this population.

"Omega-3 fatty acids have an anti-inflammatory effect, so that's one of the reasons why we suspected it may be particularly effective in obese [women](#)," Manni said.

Some epidemiological data supports the idea that omega-3s protect against breast cancer, but the findings have been inconsistent. Manni suspected that data from normal-weight women obscured the results.

Normal-weight women have less inflammation than heavier women, and are therefore less likely to benefit from anti-inflammatory omega-3s, he said.

To tease apart the effects, Manni's team, working alongside researchers from Emory University and Colorado State University, looked at the influence of prescription omega-3 supplementation on [breast density](#) in different weight women. Breast density is a well-established biomarker for [breast cancer risk](#), and may be an independent risk factor, as well. They report their results online in *Cancer Prevention Research*.

"The higher the breast density, the more likely the woman will develop breast cancer," Manni said.

The study included 266 healthy postmenopausal women with high breast density detected by routine mammograms. The women either received no treatment, the antiestrogen drug Raloxifene, the prescription omega-3 drug Lovaza or a combination of the two drugs.

At the conclusion of the two-year study, the researchers found that increasing levels of omega-3 in the blood were associated with reduced breast density—but only in women with a body mass index above 29, bordering on obesity.

Although Lovaza contains both of the fatty acids DHA—375 milligrams—and EPA—465 milligrams, only DHA blood levels were associated with breast density reduction. The researchers plan to test the effect of DHA alone in obese subjects, potentially in combination with weight loss, in a future trial.

"The finding supports the idea that omega-3s, and specifically DHA, are preferentially protective in obese postmenopausal women," Manni said. "This represents an example of a personalized approach to breast cancer prevention."

These findings may help to support future research looking at the direct effect of omega-3 supplementation on breast cancer incidence in [obese women](#).

Manni added that, with obesity-related cancers on the rise, the findings could have implications beyond breast cancer.

The researchers also made a secondary discovery. Lovaza is an omega-3 drug FDA-approved for the treatment of severe high triglycerides at the dose of 4 milligrams daily. In the current study, the combination of Lovaza and a half recommended dose of Raloxifene at 30 milligrams, was superior to the individual treatments in reducing triglycerides and

LDL—"bad"—cholesterol and increasing HDL—"good"—cholesterol.

Provided by Pennsylvania State University

Citation: Omega-3 fatty acids may lower breast cancer risk in postmenopausal obese women (2016, February 24) retrieved 5 May 2024 from <https://medicalxpress.com/news/2016-02-omega-fatty-acids-breast-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.