

Pregnancy in older age increases stroke, heart attack risk years later

February 17 2016

Women who become pregnant at age 40 or older face a greater risk of stroke and heart attack later in life than women who become pregnant at a younger age, according to research presented at the American Stroke Association's International Stroke Conference 2016.

"We already knew that older women were more likely than [younger women](#) to experience health problems during their pregnancy," said Adnan I. Qureshi, M.D., lead researcher and director of the Zeenat Qureshi Stroke Institute in St. Cloud, Minnesota. "Now, we know that the consequences of that later pregnancy stretch years into the future."

Qureshi added that this finding was especially important because more women are choosing to have children after they turn forty.

Researchers reviewed data from 72,221 women aged 50 to 79 enrolled in the Women's Health Initiative Study. Of those, 3,306 women reported pregnancies at an advanced [age](#), and researchers compared their rates of stroke, heart attack and cardiovascular death over the next 12 years with women who were pregnant at a younger age.

Researchers found that compared to pregnancy at a younger age, pregnancies at age 40 and older increased the risk of:

Ischemic stroke (clot caused), the most common type of stroke, from 2.4 percent to 3.8 percent; Hemorrhagic stroke (brain bleed), from 0.5 percent to 1 percent; Heart attack from 2.5 percent to 3 percent; and

Death from all forms of [cardiovascular disease](#) from 2.3 percent to 3.9 percent.

When researchers checked for well-known [risk factors](#) for cardiovascular disease, including high blood pressure, diabetes and high cholesterol, they found these factors explained most of the higher risk the older pregnant women faced. However, these established risk factors did not explain the link between advanced age pregnancy and stroke caused by a brain bleed. Researchers said this link needs further research. Hemorrhagic stroke accounts for about 13 percent of all strokes, according to the American Heart Association's 2016 Statistical Update.

Qureshi also noted that the only [cardiovascular risk factor](#) that did not increase with later pregnancies in the study population was smoking. Older pregnant women were less likely to smoke than younger women.

The age of last pregnancy has not been considered a risk factor for stroke, [heart attack](#) and other cardiovascular diseases. "However, women with a late pregnancy need to be aware of their increased risk and take steps to improve their cardiovascular health," he said. "And their doctors need to remain vigilant years later in monitoring these women's risk factors through physical examination and, perhaps more tests and earlier interventions to prevent [stroke](#) and other cardiovascular events."

The Women's Health Initiative Study, begun in the early 1990s by the National Institutes of Health, is one of the largest studies of women's health studies ever conducted and is highly representative of the nation's diverse population of women. Study results, he said, should be relevant to [women](#) living throughout the country.

Provided by American Heart Association

Citation: Pregnancy in older age increases stroke, heart attack risk years later (2016, February 17) retrieved 5 May 2024 from <https://medicalxpress.com/news/2016-02-pregnancy-older-age-heart-years.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.