

Which comes first: Self-reported penicillin allergy or chronic hives?

February 3 2016

People who have self-reported penicillin allergy may have a three times greater chance of suffering from chronic hives. And people who have chronic hives tend to self-report penicillin allergy at a three times greater rate than the general population. Coincidence? Authors of a new study think not.

The study, in the *Annals of Allergy, Asthma and Immunology*, the scientific publication of the American College of Allergy, Asthma and Immunology (ACAAI), examined medical records of 11,143 patients seen at the University of Pennsylvania Health System Allergy-Immunology Clinic. Of those, 220 were identified as having both self-reported penicillin allergy and chronic urticaria ([hives](#)).

"We wanted to know if there was a correlation between self-reported penicillin allergy and chronic urticaria," said lead author and [allergist](#) Susanna Silverman, MD, ACAAI member. "We found higher than expected incidence compared to the [general population](#), and we wondered if some patients who believed they had penicillin allergy might actually have chronic urticaria." Chronic urticaria is defined as hives - with or without swelling - that are present either continuously, or off and on, for longer than six weeks.

Many people have had an unfavorable response to penicillin - such as hives or swelling - at some point in their lives and are told they are allergic. Most have never seen an allergist and have never been tested for penicillin allergy. They are labeled "penicillin-allergic" on their medical

history, and are given alternative antibiotics for infections.

"It's important for anyone who thinks they have a penicillin allergy to be tested by an allergist," said allergist and study author Andrea Apter, MD, ACAAI member. "If testing finds that someone with chronic urticaria and self-reported [penicillin allergy](#) isn't allergic to penicillin, it may be that their hives are simply due to [chronic urticaria](#), or they may be more prone to rashes and hives throughout their lives, possibly due to increased skin sensitivity."

Allergists are experts in diagnosing and treating hives, and are trained to look for triggers. An allergist may recommend medications to prevent the hives or reduce the severity of symptoms. Whether the treatment is available only by prescription or over the counter will depend on several factors, including how uncomfortable the hives are making you.

Provided by American College of Allergy, Asthma, and Immunology

Citation: Which comes first: Self-reported penicillin allergy or chronic hives? (2016, February 3) retrieved 28 April 2024 from <https://medicalxpress.com/news/2016-02-self-reported-penicillin-allergy-chronic-hives.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--