

Children with autism need intervention over a long period of time

March 24 2016

9 out of 10 preschool children with autism still have major difficulties within the autism field at school age, despite having received early intervention. A majority of the parents stated that the children do not receive sufficient support at school. A new doctoral thesis at Sahlgrenska Academy explored these issues in a study where children who had received an autism diagnosis between the ages of 2 and 4.5 years of age were followed up after 8 years.

In her thesis, Martina Barnevik Olsson followed up a group of preschool children that had received a diagnosis in the autism spectrum, and who had previously received intervention from a habilitation center in Stockholm that specializes in autism, in [preschool children](#).

The parents of the children were interviewed when the children were between 9 and 13 years old.

Remaining diagnostic symptoms

A majority of the children who had been given an autism spectrum diagnosis at preschool age, had remaining diagnostic symptoms, 8 years later. They also had other neurodevelopmental disorders, even when they, as in this group, had received what is deemed to be adequate [early intervention](#).

Other problems

At follow-up, many of the children also had other difficulties that included attention and activity regulation problems, speech and language difficulties and disruptive behavior, in addition to the problems of autism.

"Most of the parents felt that the children had insufficient support in school," says Martina Barnevik Olsson.

Early regulatory problems

The thesis also investigated so-called [regulatory problems](#) (in this case problems with eating, sleeping and excessive crying) during the children's first two years of life, by mapping Children's Health Center (BVC) nurses' notes and comparing them with an age-, gender- and residence matched comparison group.

Significant difference

The results showed a [significant difference](#) between the groups, in regard to the number of times parents sought help for regulatory problems with the children. In the study group, 44 percent of the children's parents sought help at least twice for problems with food, sleep or crying, compared to only 16 percent in the comparison group. Of the 15 children with the most help-seeking occasions, 14 were from the study group.

"Even if early regulatory problems do not necessarily indicate autism, the results show the importance of noting these problems at the BVC in order to support the families and follow the children's development, as such problems can be an indicator of later developmental deviations," says Martina Barnevik Olsson.

Should be followed up

The study also showed that even those [children](#) who, over time, no longer met criteria for an autism spectrum diagnosis, should continue to be followed up, over a longer period of time. Many had remaining developmental problems and were in great need of continued support.

The thesis Autism spectrum disorders - first indicators and [school age](#) outcome was defended on March 18th.

More information: hdl.handle.net/2077/41544

Provided by University of Gothenburg

Citation: Children with autism need intervention over a long period of time (2016, March 24)
retrieved 26 April 2024 from

<https://medicalxpress.com/news/2016-03-children-autism-intervention-period.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|