

Evidence-practice gap for sealant application: Results from a dental PBRN

March 18 2016

Today at the 45th Annual Meeting & Exhibition of the American Association for Dental Research, researcher Naoki Kakudate, Kyushu Dental University, Kyushu Dental University, Japan, will present a study titled "Evidence-Practice Gap for Sealant Application: Results from a Dental PBRN." The AADR Annual Meeting is being held in conjunction with the 40th Annual Meeting of the Canadian Association for Dental Research.

In this study, the researchers aimed to examine dentist practice patterns regarding treatment recommendation of dental [sealants](#) and identify characteristics associated with this recommendation. The study was conducted using a cross-sectional questionnaire survey in Japan.

Participants were Japanese dentists (N=282) recruited from the Japanese Dental Practice Based Research Network (JDPBRN) who indicated that they do at least some restorative dentistry. Scenarios, images and questionnaire items were the same as those used in a previous U.S. DPBRN study. A series of three clinical photographs of the occlusal surface of a mandibular first molar, together with a description, were presented portraying increasing depths of cavitation. The researchers inquired about the treatment decision for each case, which had a 12-year-old patient with high caries risk. Chi-square tests were performed to assess the association between belief about the effectiveness of caries risk assessment and sealant recommendation. Multiple logistic regression analysis were conducted to evaluate the association between the decision to recommend sealants and dentist, practice and patient characteristics.

Responses were obtained from 189 dentists (67 percent). In the hypothetical scenarios, dentists' recommendations of sealants for the 12-year old patient varied from 16 percent to 26 percent. Nineteen percent of dentists recommended sealants in the absence of dark brown pigmentation. Forty-eight percent of [dentists](#) (n=91) recommended sealants to more than 25 percent of patients ages 6-18 years. Multiple logistic regression analysis suggested that the dentist's belief in the effectiveness of caries risk assessment was significantly associated with the percentage of patients who would receive sealants. Dentist practice patterns for sealant treatment recommendation vary widely. Recommending a sealant was significantly related to the dentist having a higher belief about the effectiveness of caries risk assessment.

This research was supported by NIH grants U01-DE-16746, U01-DE-16747 and U19-DE-22516.

More information: This is a summary of oral presentation #0982, "Evidence-Practice Gap for Sealant Application: Results from a Dental PBRN," which will be presented on Friday, March 18, 2016, 10:45 a.m. - 11 a.m. at the Los Angeles Convention Center, room #406A.

Provided by International & American Associations for Dental Research

Citation: Evidence-practice gap for sealant application: Results from a dental PBRN (2016, March 18) retrieved 3 May 2024 from <https://medicalxpress.com/news/2016-03-evidence-practice-gap-sealant-application-results.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
