


For prostate cancer, more radiation may not improve survival

March 29 2016



Micrograph showing prostatic acinar adenocarcinoma (the most common form of prostate cancer) Credit: Wikipedia, [CC BY-SA 3.0](https://creativecommons.org/licenses/by-sa/3.0/)

New technology has enabled doctors to administer higher doses of radiation to prostate cancer patients with fewer side effects. However, a new study shows that escalating the dose may not actually help a patient in the long term, at least not patients with localized prostate cancer. The results were published online last week in the *American Journal of Clinical Oncology*.

"In the field of [radiation oncology](#), we often assume that the highest dose that the body can tolerate will be most effective at killing cancer," says Dr. Robert Den, M.D., a researcher at the Sidney Kimmel Cancer Center at Thomas Jefferson University and senior author on the paper. "Our results argue that this may not be the case, at least not with lower-risk [prostate cancer](#) patients."

Dr. Den, an Associate Professor of Radiation Oncology, Cancer Biology, and Urology at Jefferson, and colleagues analyzed data from 12 [randomized controlled trials](#) of external beam radiation treatment for men with non-metastatic prostate cancer, which included a total of 6884 patients. By pooling data from multiple clinical trials, the researchers were able to see trends that would not have been apparent in the individual studies.

Rather than use the typical proxy for patient improvement, the prostate cancer antigen (PSA) test, the researchers looked at long term outcomes such as the development of metastatic cancer and death from cancer. They found that while PSA levels decreased as patients received higher doses of radiation, the overall survival and incidence of metastases, among other measures, did not improve.

"It's important to check our assumptions," says Adam Dicker, Chair of Radiation Oncology at the Sidney Kimmel Medical College of Thomas Jefferson University. "This study suggests that our reliance on the PSA test as a proxy for patient outcomes may not as useful as many researchers thought, which has broad implications for the design of future clinical trials and the interpretation of current and previous studies."

Dr. Den's study also demonstrated that increasing dose was not associated with worse treatment toxicity, suggesting that current practices are safe. "These data suggest that other therapies may be

needed with radiation to increase survival," says Dr. Den.

More information: Nicholas G. Zaorsky et al. Impact of Radiation Therapy Dose Escalation on Prostate Cancer Outcomes and Toxicities, *American Journal of Clinical Oncology* (2016). [DOI: 10.1097/COC.0000000000000285](https://doi.org/10.1097/COC.0000000000000285)

Provided by Thomas Jefferson University

Citation: For prostate cancer, more radiation may not improve survival (2016, March 29) retrieved 30 March 2023 from <https://medicalxpress.com/news/2016-03-prostate-cancer-survival.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.