

## Study reports significant link between nightmares and suicidal behavior

March 16 2016

---



Credit: Vera Kratochvil/public domain

A new study is the first to report that the relationship between

nightmares and suicidal behaviors is partially mediated by a multi-step pathway via defeat, entrapment and hopelessness.

Results show that [suicidal thoughts](#), plans or attempts were present in 62 percent of participants who experienced nightmares and only 20 percent of those without nightmares. Multiple analyses suggest that nightmares may act as a stressor in people with post-traumatic stress disorder (PTSD). The nightmares may trigger specific types of negative cognitive thoughts—such as defeat, entrapment and hopelessness—which reinforce suicidal thoughts and behaviors. The pathways between nightmares and suicidal behaviors appear to operate independent of comorbid insomnia and depression.

"PTSD increases the risk of suicidal thoughts and behavior, and our study shows that nightmares, a hallmark symptom of PTSD, may be an important treatment target to reduce suicide risk," said principal investigator Donna L. Littlewood, PhD researcher in medical and human sciences at The University of Manchester. "This study emphasizes the importance of specifically assessing and targeting nightmares within those individuals experiencing PTSD. In addition, monitoring and targeting levels of negative cognitive appraisals such as defeat, entrapment, and hopelessness, may reduce suicidal thoughts and behaviors."

Study results are published in the March 15, 2016 issue of the *Journal of Clinical Sleep Medicine*.

The American Academy of Sleep Medicine reports that nightmares are vivid, realistic and disturbing dreams typically involving threats to survival or security, which often evoke emotions of anxiety, fear or terror. A nightmare disorder may occur when repeated nightmares cause distress or impairment in social or occupational functioning. Nightmares beginning within three months of a trauma are present in up to 80

percent of patients with PTSD, and these post-traumatic nightmares may persist throughout life.

Data for this study was collected from 91 participants who had experienced traumatic events, 51 of whom met criteria for PTSD currently, and an additional 24 of whom reported a prior diagnosis of PTSD. Nightmares were measured by summing the frequency and intensity ratings of relevant items on the clinician-administered PTSD scale. Participants also completed questionnaire measures of [suicidal behavior](#), hopelessness, defeat, and entrapment. Given the interactions between insomnia, PTSD, and suicide, a measure of insomnia was included as a covariate. Analysis was also conducted with and without those participants who had comorbid depression.

This study was conducted under the supervision of Simon D. Kyle, PhD, of the Sleep and Circadian Neuroscience Institute at the University of Oxford in the U.K.

The authors suggest there are additional pathways underpinning the relationship between [nightmares](#) and suicide that should be identified through further research.

**More information:** "Nightmares and Suicide in Posttraumatic Stress Disorder: The mediating Role of Defeat, Entrapment, and Hopelessness", *Journal of Clinical Sleep Medicine*, 2016.

Provided by American Academy of Sleep Medicine

Citation: Study reports significant link between nightmares and suicidal behavior (2016, March 16) retrieved 23 April 2024 from <https://medicalxpress.com/news/2016-03-significant-link-nightmares-suicidal-behavior.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.