

Teens with autism more likely to develop type 2 diabetes

March 28 2016



(HealthDay)—Adolescents and young adults with autism spectrum

disorder (ASD) are more likely to develop type 2 diabetes mellitus (T2DM), according to a study published online March 22 in *Diabetes Care*.

Mu-Hong Chen, M.D., from the Taipei Veterans General Hospital in Taiwan, and colleagues used the Taiwan National Health Insurance Research Database to identify 6,122 adolescents and young adults with ASD and 24,488 age- and sex-matched control subjects (2002 to 2009). Patients were monitored until the end of 2011 to identify those who developed T2DM.

The researchers found that adolescents (hazard ratio [HR], 2.71) and [young adults](#) (HR, 5.31) with ASD had a higher risk of developing T2DM versus those without ASD, after adjusting for demographic data, use of atypical antipsychotic, and medical comorbidities. There was also a higher likelihood of subsequent T2DM tied to short-term (HR, 1.97) and long-term (HR, 1.64) use of atypical antipsychotics.

"Further research is necessary to investigate the common pathophysiology of ASD and T2DM," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Teens with autism more likely to develop type 2 diabetes (2016, March 28) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-03-teens-autism-diabetes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--