

## Study finds adolescent tobacco users commonly report light smoking

April 30 2016

---

A new research abstract being presented at the Pediatric Academic Societies (PAS) 2016 Meeting reveals new details about teen smoking. Most young smokers report that they don't light up every day, and many smoke only a few cigarettes on the days they do smoke. These teens are less likely to identify as smokers, even as they face health risks comparable to heavier tobacco use.

The study, "Intermittent and Light Smoking Patterns among Youth," followed 602 young smokers enrolled in 115 pediatric practices in 39 states from the AAP Pediatric Research in Office Settings (PROS) practice based research network. Teens completed a baseline survey and follow-up surveys at 4-6 weeks, 6 months and 1 year. Researchers said 73 percent of the teens reported being intermittent smokers, and 69 percent reported [smoking](#) fewer than five cigarettes per day. The intermittent and light smokers, more likely to be younger, female and above-average students, were less likely than heavier smokers to identify as smokers.

"These intermittent and light patterns of smoking are important for clinicians to understand, because it impacts the way we talk to teens about smoking," said presenting author Julie Gorzkowski, MSW, LSW, senior research associate with the Julius B. Richmond Center of Excellence at the American Academy of Pediatrics. "Asking a [teen](#) if they're a smoker may not get you the information you need, because a teen might identify as a non-smoker but still smoke on the weekends or at parties. We need to tailor the language we use to the behavior patterns

that are more relevant for them."

Of particular concern, the researchers said, is that light and intermittent [smokers](#) reported that they were less likely to want to quit or to make attempts to quit.

"Teens who use cigarettes less often are still subject to the dangerous health effects of smoking," said co-author Jonathan Klein, MD, MPH, FAAP, founding director of the AAP Richmond Center. He added that developing adolescent brains are particularly sensitive to nicotine and even a few instances of smoking a cigarette early in life make a teen more likely to become addicted and to continue smoking down the road.

"It's important to understand that there is no safe level of tobacco use," Dr. Klein said. "Experimenting with tobacco shouldn't be considered a normal part of adolescence. Clinicians should screen for [light](#) and intermittent smoking with teen patients, and work to help these patients stop smoking now."

**More information:** [www.abstracts2view.com/pas/view.asp?nu=PAS16L1\\_3115.3](http://www.abstracts2view.com/pas/view.asp?nu=PAS16L1_3115.3)

Provided by American Academy of Pediatrics

Citation: Study finds adolescent tobacco users commonly report light smoking (2016, April 30) retrieved 24 April 2024 from <https://medicalxpress.com/news/2016-04-adolescent-tobacco-users-commonly.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--