

Increased BMI during adolescence predicts fatal cardiovascular events in adulthood

April 14 2016, by Dov Smith



Credit: Hebrew University of Jerusalem

Overweight and obesity in adolescents have increased substantially in recent decades, and currently affect a third of the adolescent population

in some developed countries. This is an important public health concern because obesity early in life is considered to be a risk factor for death from cardiovascular disease and from all causes in adulthood.

Some studies suggest that an elevated body-mass index is associated with an [increased risk](#) of death from cardiovascular causes. However, a determination of the BMI threshold that is associated with increased risk of fatality remains uncertain. (BMI is a calculation of a person's weight in kilograms divided by the square of their height in meters, to quantify body mass and enable categorization as underweight, normal weight, overweight, or obese.)

In light of the worldwide increase in [childhood obesity](#), Prof. Jeremy Kark and Dr. Hagai Levine from the Hebrew University-Hadassah Braun School of Public Health and Community Medicine, in the Hebrew University of Jerusalem's Faculty of Medicine, together with Dr. Gilad Twig of Sheba Medical Center and other colleagues in Israel, set out to determine the association between body-mass index (BMI) in late adolescence and death from cardiovascular causes in adulthood.

Their study, which appears in the *New England Journal of Medicine*, was based on a national database of 2.3 million Israeli 17 year olds in whom height and weight were measured between 1967 and 2010. The researchers assessed the association between BMI in late adolescence and death from [coronary heart disease](#), stroke, and sudden death in adulthood by mid-2011.

During 42,297,007 person-years of follow-up, 2918 of 32,127 deaths (9.1%) were from cardiovascular causes, including 1497 from [coronary heart disease](#), 528 from stroke, and 893 from sudden death.

The results showed an increase in the risk of cardiovascular death in the group that was considered within the "accepted normal" range of BMI,

in the 50th to 74th percentiles, and of death from coronary heart disease at BMI values above 20. The researchers concluded that even BMI considered "normal" during adolescence was associated with a graded increase in cardiovascular and all-cause mortality during the 40 years of follow-up. This included increased rates of death from coronary heart disease, stroke, and total cardiovascular causes among participants.

As BMI scores increased into the 75th to 84th percentiles, adolescent obesity was associated with elevated risk of death from coronary heart disease, stroke, sudden death from unknown causes, and death from total cardiovascular causes, as well as death from non-cardiovascular causes and death from all causes. Participants also had an increased risk of sudden death.

The rates of [death](#) per person-year were generally lowest in the group that had BMI values during adolescence in the 25th to 49th percentiles, although higher rates were observed among those below the 5th percentile.

How might adolescent BMI influence cardiovascular outcomes in adulthood? The researchers considered two possible pathways. First, obesity may be harmful during adolescence, since it has been associated with unfavorable metabolic abnormalities through risk factors such as unfavorable plasma lipid or lipoprotein levels, increased blood pressure, impaired glucose metabolism, insulin resistance, and formation of coronary and aortic atherosclerotic plaques. Furthermore, the timing of exposure to obesity during a person's lifetime may play an important role. Second, BMI tends to "track" along the life course so that overweight adolescents tend to become overweight or obese adults, and overweight or obesity in adulthood affects the risk of cardiovascular disease.

"Our findings appear to provide a link between the trends in adolescent

overweight during the past decades and coronary mortality in midlife," said the paper's senior author, Prof. Jeremy Kark of the Hebrew University-Hadassah Braun School of Public Health and Community Medicine. "The continuing increase in adolescent BMI, and the rising prevalence of overweight and obesity among adolescents, may account for a substantial and growing future burden of [cardiovascular disease](#), particularly coronary heart disease."

More information: Gilad Twig et al. Body-Mass Index in 2.3 Million Adolescents and Cardiovascular Death in Adulthood, *New England Journal of Medicine* (2016). [DOI: 10.1056/NEJMoa1503840](https://doi.org/10.1056/NEJMoa1503840)

Provided by Hebrew University of Jerusalem

Citation: Increased BMI during adolescence predicts fatal cardiovascular events in adulthood (2016, April 14) retrieved 2 May 2024 from <https://medicalxpress.com/news/2016-04-bmi-adolescence-fatal-cardiovascular-events.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
