

Childhood asthma overdiagnosed, argue leading respiratory doctors

April 5 2016



Children with asthma use inhalers to relieve some of their symptoms, which include coughing, wheezing, chest tightness and shortness of breath. Credit: Tradimus / Wikimedia commons / CC BY-SA 3.0

Doctors are overdiagnosing asthma, with inhalers frequently dispensed for no good reason, to the point that they have "almost become a fashion accessory," argue two leading respiratory doctors in a leading article published online in the *Archives of Disease in Childhood*.



In the past <u>asthma</u> was undoubtedly underdiagnosed, but the evidence now is that the pendulum has swung too far in the opposite direction, insist Professor Andrew Bush and Dr Louise Fleming of Imperial College and Royal Brompton & Harefield NHS Foundation Trust, in the journal.

It matters, not only because of the cost of inhalers, but also because of their side effects, which are more likely to occur in patients who are prescribed an inappropriate dose, they say.

"Inhaled corticosteroids, when properly used, dramatically improve quality of life and reduce the risk of asthma attacks and mortality," they write.

But there are potential side effects associated with their use, they point out. These include growth suppression and the dampening down of immune cell activity in the airways and the subsequent heightened risk of respiratory infections.

"There is also evidence that systemic absorption of [inhaled corticosteroids] depends not just on the prescribed dose, but is greater if the dose is inappropriately high for the degree of airway inflammation," they caution.

Key to diagnosing asthma correctly is a detailed knowledge of normal respiratory symptoms in healthy children, they say. But as the National Review of Asthma deaths shows, children who have asthma still die because of failures in basic management, including not taking inhalers that they really need.

"We propose that one contributing factor is that the diagnosis of asthma has been trivialised and inhalers dispensed for no good reason, and have become almost a fashion accessory," they write.



"The result is, the fact that asthma is a killing disease if not correctly managed, is overlooked," they say.

All too often, the hallmark symptom of asthma—wheeze—is used imprecisely by both parents and healthcare professionals. More objective evidence, such as simple breathing tests, is required before a diagnosis is made, they suggest.

"Is there any other chronic disease in the world in which children are committed to potentially hazardous, long term therapy without every effort being made objectively to document the diagnosis?" they ask.

They outline various approaches to aid correct diagnosis, including the need to remember that many children outgrow asthma symptoms, and that treatment shouldn't simply be stepped up if the child fails to respond, because there's a chance the diagnosis might not be right in the first place.

More information: Is asthma overdiagnosed? <u>DOI:</u> 10.1136/archdischild-2015-309053

Provided by British Medical Journal

Citation: Childhood asthma overdiagnosed, argue leading respiratory doctors (2016, April 5) retrieved 3 May 2024 from

https://medicalxpress.com/news/2016-04-childhood-asthma-overdiagnosed-respiratory-doctors.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.