

Combination therapy for acute lymphoblastic leukemia

April 7 2016

A subset of acute lymphoblastic leukemia (ALL) patients is positive for the Philadelphia chromosome, which is generated by a specific translocation event that causes a fusion between the *BCR* and *ABL1* genes. These patients unfortunately have a poorer prognosis and a more limited response to tyrosine kinase inhibitors that are used to treat other forms of ALL.

In this issue of *JCI Insight*, Charles Mulligan of St. Jude Children's Research Hospital and colleagues show that a combination approach using [tyrosine kinase inhibitors](#) and an inhibitor of focal adhesion kinase (FAK) was much more effective in preclinical models of BCR-ABL1⁺ B-progenitor ALL. They demonstrate that FAK is overexpressed in several models of BCR-ABL1⁺ ALL and that the FAK inhibitor VS-4718 synergizes with tyrosine inhibitor dasatinib to decrease tumor growth in mouse models.

Collectively, their work indicates that a combination approach, targeting both FAK and [tyrosine kinases](#), is a promising treatment strategy for this subset of ALL.

More information: Michelle L. Churchman et al. Synergism of FAK and tyrosine kinase inhibition in Ph⁺ B-ALL, *JCI Insight* (2016). [DOI: 10.1172/jci.insight.86082](https://doi.org/10.1172/jci.insight.86082)

Provided by Journal of Clinical Investigation

Citation: Combination therapy for acute lymphoblastic leukemia (2016, April 7) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2016-04-combination-therapy-acute-lymphoblastic-leukemia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.