

Diffusion-weighted imaging + MRI OK for undescended testes

April 20 2016



(HealthDay)—Combined diffusion-weighted imaging (DWI) and

magnetic resonance imaging (MRI) show a greater performance compared to conventional MRI alone for identification of non-palpable undescended testes (UDTs), according to a study published online April 6 in the *Journal of Medical Imaging and Radiation Oncology*.

Sally Emad-Eldin, M.D., from Cairo University Hospitals, and colleagues evaluated the diagnostic performance of combined DWI and conventional MRI, including fat-suppression T2WI, for identification and localization of non-palpable UDTs in 40 consecutive patients with 47 non-palpable undescended testes (33 unilateral cases and seven bilateral cases; mean age 7.5 years).

The researchers found that the final diagnoses of the location of UDTs based on laparoscopy findings were: intra-canalicular (18 cases), low intra-abdominal (six cases), and high intra-abdominal (five cases). There were 18 cases of absent or vanishing testes. Combined DWI and conventional MRI had a diagnostic accuracy, sensitivity, and specificity of 95.7, 93.5, and 100 percent, respectively.

"Based on our findings, we can obviate the need for diagnostic laparoscopy in patients who had preoperative detection of inguinal testes or nubbins," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Diffusion-weighted imaging + MRI OK for undescended testes (2016, April 20)
retrieved 25 April 2024 from
<https://medicalxpress.com/news/2016-04-diffusion-weighted-imaging-mri-undescended.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.