

First implantable hemodynamic monitoring device in single ventricle Fontan anatomy

April 28 2016

While the Fontan procedure has improved the short- and mid-term outcomes for patients born with single ventricle anatomy, long-term complications of Fontan circulation include heart failure. These complications are thought to be secondary to elevated central venous pressure, chronic venous congestion and low cardiac output.

In the recent case series published in the journal *Catheterization and Cardiovascular Interventions*, clinicians from The Heart Center at Nationwide Children's Hospital and The Ohio State University Wexner Medical Center describe the first implantable hemodynamic monitor (IHM) placement in single ventricle Fontan anatomy. To date, Nationwide Children's is the first pediatric hospital to perform the procedure, while Ohio State's Wexner Medical Center is the first hospital to perform the procedure in a Fontan patient.

"This pressure gradient between central venous pressure and a combination of ventricular end diastolic pressure and resistance of flow across the pulmonary vascular bed is critical to cardiac output," explains Darren Berman, MD, co-director of Cardiac Catheterization and Interventional Therapy in The Heart Center at Nationwide Children's. "Recently, implantable hemodynamic monitors have been used to monitor pressures in patients with left heart disease."

Noninvasive studies are unable to determine these pressures in the Fontan anatomy, thus limiting physicians' understanding of the changes in pressure for a Fontan patient in <u>heart failure</u>, according to Curt



Daniels, MD, director of Adolescent and Adult Congenital Heart Disease in The Heart Center at Nationwide Children's and professor of Clinical Cardiology at The Ohio State University College of Medicine.

"Our ultimate goal is to improve care for patients with <u>congenital heart</u> <u>disease</u>," says Dr. Daniels. "We are excited to be leading the charge to incorporate the use of this tool to learn more about the hemodynamic changes in real life ambulatory situations in Fontan patients."

Two patients with single ventricle anatomy and Fontan physiology, both with New York Heart Association (NYHA) functional class III <u>heart</u> failure, were implanted with the CardioMEMS HF System by Drs. Berman and Daniels. Real-time pulmonary artery catheter tracings were shown to correlate with IHM tracings in Fontan anatomy. Additionally, home transmission of IHM readings was successful.

"We have shown early safety and feasibility of placement of the IHM in these two cases, and we will follow these patients to monitor the safety in the medium- and long-term, including assessment for pulmonary arterial thrombus," says Dr. Berman, who is also an assistant professor at OSUCM.

More information: Elisa A. Bradley et al, First implantable hemodynamic monitoring device placement in single ventricle fontan anatomy, *Catheterization and Cardiovascular Interventions* (2016). DOI: <u>10.1002/ccd.26498</u>

Provided by Nationwide Children's Hospital

Citation: First implantable hemodynamic monitoring device in single ventricle Fontan anatomy (2016, April 28) retrieved 28 April 2024 from



https://medicalxpress.com/news/2016-04-implantable-hemodynamic-device-ventricle-fontan.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.