

Risk of liver cancer from hepatitis B persists even after clearing the virus

April 22 2016

Long-term infection with hepatitis B virus (HBV) can cause liver inflammation and increase the risk of liver cancer. Researchers from the US Centers for Disease Control and Prevention, however, found that resolving HBV infection was not associated with reduced rates of liver cancer.

A total of 238 patients who resolved HBV infection and 435 patients who did not were selected from among 1346 patients with chronic HBV infection followed for up to 31 years. The [liver cancer](#) risk, measured as a hazard ratio (HR), in patients after resolving HBV infection was similar compared with the risk in patients who did not resolve infection (HR: 0.7; 95% confidence interval: 0.2-2.4).

"Since the risk of liver cancer persists among adults with apparent cure of the infection, they might still need to be followed closely," said Dr. Prabhu Gounder, lead author of the *Alimentary Pharmacology & Therapeutics* article.

More information: P. P. Gounder et al, Nested case-control study: hepatocellular carcinoma risk after hepatitis B surface antigen seroclearance, *Alimentary Pharmacology & Therapeutics* (2016). [DOI: 10.1111/apt.13621](#)

Provided by Wiley

Citation: Risk of liver cancer from hepatitis B persists even after clearing the virus (2016, April 22) retrieved 1 May 2024 from <https://medicalxpress.com/news/2016-04-liver-cancer-hepatitis-persists-virus.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.