

Psoriasis severity linked to rising risk of abdominal aortic aneurysms

April 14 2016

Psoriasis sufferers may face a higher risk of developing abdominal aortic aneurysms, according to new research in *Arteriosclerosis, Thrombosis and Vascular Biology*, an American Heart Association journal.

The study also found that the more severe the [psoriasis](#), the greater the risk for abdominal aortic aneurysms (AAA).

Psoriasis is a chronic autoimmune disease—a condition in which the body's immune system attacks healthy cells—characterized by patches of abnormal skin. These skin patches are typically red, itchy and scaly. AAA is an enlargement of the main blood vessel that supplies blood to the abdomen, and usually causes no symptoms until it ruptures, which could prove fatal.

"Psoriasis must be considered as a systemic inflammatory disease rather than an isolated skin disease. Increased awareness on heightened risk of other cardiovascular diseases, including AAA, in patients with psoriasis is also required," said Usman Khalid, M.D., lead author and Ph.D. fellow in the Department of Cardiology, at Herlev and Gentofte Hospital in Denmark.

Using Danish nationwide registers, researchers identified 59,423 patients with mild psoriasis and 11,566 patients with [severe psoriasis](#) between 1997 and 2011. These patients were followed until diagnosis of abdominal aortic aneurysms, death, migration or the study end date. The

rate of developing [abdominal aortic aneurysms](#) (per 10,000 person-years) were:

3.72 for general population, 7.30 for patients with mild psoriasis, and 9.87 for patients with severe psoriasis.

"Compared to the general population, the adjusted incidence rate ratios were significantly increased for severe psoriasis at 1.67. That's a 67 percent greater risk of AAA likelihood for severe psoriasis sufferers," Khalid said. "Clinicians need to educate and assist their patients with psoriasis in lifestyle and [risk factor modification](#) to facilitate [cardiovascular disease risk](#) reduction."

Khalid said their study adds significantly to current evidence of psoriasis as a clinically relevant risk factor for cardiovascular disease and may require further investigation to answer questions, such as whether patients with psoriasis should undergo increased ultrasonic screening for AAA, and whether anti-inflammatory treatment of psoriasis may reduce the risk of AAA.

Provided by American Heart Association

Citation: Psoriasis severity linked to rising risk of abdominal aortic aneurysms (2016, April 14) retrieved 17 April 2024 from <https://medicalxpress.com/news/2016-04-psoriasis-severity-linked-abdominal-aortic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--