


Study supports single-question alcohol screen for adolescents

April 7 2016



“Alcohol Screening and Brief Intervention for Youth: A Practitioner’s Guide” is designed to help health care professionals quickly identify youth at risk for

alcohol-related problems.

A single screening question about drinking frequency in the past year could help doctors identify adolescents at risk for alcohol problems, according to a new study funded by the National Institute on Alcohol Abuse and Alcoholism (NIAAA), part of the National Institutes of Health. Conducted by researchers at the University of Pittsburgh, who collaborated with a network of rural primary care practitioners, the study also supports the use of the age-based screening thresholds put forward in NIAAA's [Alcohol Screening and Brief Intervention for Youth: A Practitioner's Guide](#).

"Primary care physicians are encouraged to screen adolescents for [alcohol problems](#), yet many do not, citing time constraints and other issues," said NIAAA Director George Koob, PhD. "This study demonstrates that simple [screening](#) tools such as those in NIAAA's Youth Guide are efficient and effective."

Researchers led by Duncan B. Clark, M.D., Ph.D., Professor of Psychiatry at the University of Pittsburgh Medical Center, asked nearly 1,200 young people ages 12 through 20 at [primary care](#) clinics in rural Pennsylvania about their alcohol use and screened them for [alcohol use disorder \(AUD\)](#) using a computer-based questionnaire. They found that 10 percent of rural youth over age 14 met the diagnostic criteria for AUD in the past year, as specified in the Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-5). The new findings are now online in *The Journal of Pediatrics*.

Adolescents between 12 and 17 years of age who reported drinking at least one standard drink on three or more days in the past year were most at risk for alcohol problems. Among those who drank at this level, 44

percent were found to have AUD. Screening for AUD based on the 3-day guideline demonstrated 91 percent sensitivity—indicating that a youth with AUD was likely to be detected by the screen—and 93 percent specificity—meaning those without AUD were likely have a negative result. A negative screen (fewer than three drinking days in the past year) effectively ruled out AUD, with 99 percent not having the disorder.

For adolescents ages 18-20, the researchers found that the best screen for alcohol problems was to ask whether individuals had engaged in 12 or more drinking days (i.e., a day in which at least one standard drink is consumed) in the past year. Thirty-one percent of those who reported drinking at this level had AUD.

"This finding confirms that a single question can be an effective screen for AUD," said Dr. Clark. "We found that this information could be readily collected through our tablet computer system in busy rural clinic settings."

Dr. Clark and his colleagues also assessed screening methods outlined in the NIAAA Youth Screening Guide, which uses age-specific alcohol frequency questions to identify moderate and highest levels of AUD risk. "Moderate risk" is defined as one or more drinking days in the past year for ages 12-15, three or more days per year for ages 16-17, and 12 or more days per year for 18-year-olds. "Highest risk" is defined as three or more days per year for youth ages 12-15, 12 or more days for 16-year-olds, 24 or more days for 17-year-olds, and 52 or more days for 18-year-olds.

Researchers found both sets of NIAAA guidelines to be effective screens for AUD. The moderate risk guidelines showed 85 percent sensitivity and 87 percent specificity, while the highest risk guidelines showed 91 percent sensitivity and 93 percent specificity.

The researchers note that, as national studies document higher rates of alcohol use among rural [youth](#), identifying problem drinking is of particular importance for primary care providers in rural settings. They conclude that an [alcohol](#) use frequency screen followed by a diagnostic evaluation for those who screen positive would be "a simple, brief, and cost-effective clinical assessment procedure."

More information: The NIAAA Alcohol Screening and Brief Intervention for Youth: A Practitioner's Guide is available at niaaa.nih.gov/publications/clinician-intervention-youth

Duncan B. Clark et al. Screening for Underage Drinking and Diagnostic and Statistical Manual of Mental Disorders, 5th Edition Alcohol Use Disorder in Rural Primary Care Practice, *The Journal of Pediatrics* (2016). [DOI: 10.1016/j.jpeds.2016.02.047](https://doi.org/10.1016/j.jpeds.2016.02.047)

Provided by National Institutes of Health

Citation: Study supports single-question alcohol screen for adolescents (2016, April 7) retrieved 4 May 2024 from

<https://medicalxpress.com/news/2016-04-single-question-alcohol-screen-adolescents.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.