

New model for active surveillance of prostate cancer tested

April 27 2016



Christopher Kane, M.D., is chair of the Department of Urology at UC San Diego Health. Credit: UC San Diego Health

Urologists at University of California, San Diego School of Medicine and Genesis Healthcare Partners have tested a new model of care for patients with low-risk prostate cancer. The evidence-based approach uses best practices to appropriately select and follow patients to avoid disease overtreatment. Results of the three-year study are now published online in the journal of *Urology*.

"Active surveillance is a strategy that is recommended by physician and quality organizations to avoid the overtreatment of slow-growing [prostate cancer](#)," said Christopher Kane, MD, senior author and chair of the Department of Urology at UC San Diego Health. "Acceptance of this strategy by [patients](#) and urologists, however, has lagged for a number of reasons. What we have developed is a safe method to enhance acceptance and use of this disease management approach."

Active surveillance (AS) is the practice of closely monitoring slow, indolent forms of prostate cancer with prostate-specific antigen (PSA) blood tests, digital rectal prostate exams and, potentially, biopsies. Kane added that AS recognizes that there is a large group of men with a form of low-grade prostate cancer whose long-term survival is not impacted by non-treatment.

"With this [new model](#), we were able to increase rates of surveillance to benefit patients through use of provider education and a standardized report card," said Franklin Gaylis, MD, Chief Scientific Officer, Genesis Healthcare and the study's first author. "With this university and private-practice research collaboration, we were able monitor 190 patients undergoing [active surveillance](#) while evaluating the effectiveness of our own individual clinical practices."

Urologists at Genesis Healthcare and UC San Diego Health jointly developed a reporting mechanism to improve the process of tracking patients with prostate cancer. The research team developed standardized selection criteria based on scientific literature for patients to be followed with AS according to tumor characteristics, including clinical cancer staging, Gleason (pathology grading) and PSA scores. In addition, comparative dashboards were developed to show individual physician AS adoption rates compared to their peers.

"What we found is that active surveillance increased from 43.75 percent

to 82.6 percent among the very low-risk patients," said Gaylis, a voluntary professor in the Division of Urology at UC San Diego School of Medicine. "Besides this approach, enhancing quality of care through established standardized processes and outcomes feedback, there may be a benefit from a cost-savings perspective. This model may be particularly helpful as the U.S. switches from a volume to value-based system of care for reimbursement requiring physicians to improve quality while at the same time reducing cost."

Prostate cancer is the most prevalent solid organ malignancy among American men, accounting for almost 30 percent of new cancer diagnoses. The National Institutes of Health estimates approximately 220,800 new cases of prostate cancer will be identified in 2015.

Provided by University of California - San Diego

Citation: New model for active surveillance of prostate cancer tested (2016, April 27) retrieved 28 April 2024 from <https://medicalxpress.com/news/2016-04-surveillance-prostate-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.