

ATS: Reintubation more likely for nighttime extubations

May 18 2016



(HealthDay)—Patients extubated at nighttime are more likely to be

reintubated and have higher mortality than those extubated during the day, according to a study presented at the annual meeting of the American Thoracic Society, held from May 13 to 18 in San Francisco.

Hayley B. Gershengorn, M.D., from the Albert Einstein College of Medicine in New York City, and colleagues performed a [retrospective cohort study](#) of 32,760 mechanically ventilated (MV) adults aged 18 years and older admitted to 76 U.S. intensive care units (ICUs). The authors examined the frequency of and patient characteristics associated with nighttime extubations across ICUs.

The researchers found that 18.2 percent of patients were extubated overnight, with a median rate of nighttime extubations of 15.2 percent. After multivariable adjustment, patient factors associated with nighttime extubation included diagnosis of cardiovascular disease, sepsis, trauma, or neurologic (versus respiratory) disease; higher acute physiology score; absence of MV on ICU admission; and MV duration

Citation: ATS: Reintubation more likely for nighttime extubations (2016, May 18) retrieved 8 May 2024 from <https://medicalxpress.com/news/2016-05-ats-reintubation-nighttime-extubations.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--