

Children with ADHD sleep both poorly and less

May 4 2016

A new study from Aarhus University has now documented that there is some truth to the claim by parents of children with ADHD that their children have more difficulty falling asleep and that they sleep more poorly than other children.

Studies have shown that up to seventy per cent of parents of children with ADHD report that the children have difficulty falling asleep and that they spend a long time putting them to bed. However, scientific studies that measure [sleep quality](#) using electrodes have so far failed to demonstrate a correlation between [sleep](#) quality and ADHD. But a new Danish study now shows that children with ADHD actually do sleep worse than other children:

"Our study will confirm what many parents have experienced, which is that children with ADHD take longer to fall asleep at night. With our measurements we can also see that these children experience more disturbed sleep including less deep sleep. If you only look at length of sleep, children in the ADHD group sleep for 45 minutes less than children in the [control group](#)," says PhD from Aarhus University and medical doctor at the Child and Adolescent Psychiatric Hospital, Risskov, Anne Virring Sørensen, who is behind the PhD study.

Two out of three children with ADHD have one or more additional psychiatric diagnoses in addition to ADHD, which probably increases the risk of sleep disturbance. But even when the researchers look at the children who have only been diagnosed with ADHD, they see a big

difference in the sleep patterns of the control group and the ADHD group.

Opposite sleep patterns during the day

The researchers also studied sleep patterns during the day. The findings surprised the researchers.

"Unlike in the evening we could see that there was a tendency for the children with ADHD to fall asleep faster during the day than the children in the control group. This is somewhat surprising when you take into account that ADHD is associated with characteristics such as hyperactivity. But this hyperactivity could be compensatory behaviour for not being able to doze off during the day" says Anne Virring Sørensen.

The fact that researchers have not previously been able to demonstrate a correlation between ADHD and poorer sleep could be due to different measuring methods.

"In our study the children had electrodes attached to their heads for what is known as a polysomnography at the hospital in the afternoon, but they slept in their familiar home surroundings. In previous studies children have been admitted to specialist sleep centres at hospital to measure sleep via a polysomnographic study," says Anne Virring Sørensen.

Many children with ADHD are currently given medicine to help them fall asleep. Anne Virring Sørensen emphasises that none of the children received medicine while taking part in the study.

Important study

She believes that the study is important in both the short and long term:

"I think many parents and clinicians are very pleased to receive confirmation that poor [sleep patterns](#) can now be demonstrated and that there is probably a correlation with the ADHD diagnosis. The next step is, of course, to find out where this correlation lies so we can develop better treatments in the long term. Our survey is an important foundation for further studies," she says.

The study was recently published in the *Journal of Sleep Research*.

Facts about the study

- A total of 76 children with ADHD and an average age of 9.6 years participated in the study.
- The control group consisted of 25 healthy children.
- Two different types of study were carried out:
- Outpatient sleep examinations with electrodes during the night (polysomnography).
- Multiple sleep latency tests which measured how quickly the children fell asleep (four times twenty minutes on the same day).
- The study is the largest so far to include both research methods and to include both [children](#) with and without an ADHD diagnosis.
- The study was carried out by researchers from Aarhus University, Aarhus University Hospital Risskov, Rigshospitalet and the University of Copenhagen.

More information: Anne Virring et al, Disturbed sleep in attention-deficit hyperactivity disorder (ADHD) is not a question of psychiatric comorbidity or ADHD presentation, *Journal of Sleep Research* (2016).

[DOI: 10.1111/jsr.12377](https://doi.org/10.1111/jsr.12377)

Provided by Aarhus University

Citation: Children with ADHD sleep both poorly and less (2016, May 4) retrieved 25 April 2024 from <https://medicalxpress.com/news/2016-05-children-adhd-poorly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.