

# Study of elite paralympic athletes supports benefits of exercise for children with cerebral palsy

May 6 2016

---

For highly trained Paralympic athletes with cerebral palsy (CP), bone mineral density and other measures of body composition are similar to those of able-bodied adults of similar age, reports a study in the *American Journal of Physical Medicine & Rehabilitation*.

While elite-level [athletes](#) with hemiplegic CP still have reduced muscle mass on the side of their body affected by neuromuscular impairment, physical training can offset other CP-related changes in body composition and physiology, suggests the study by Phoebe Runciman, PhD, of University of Cape Town, South Africa, and colleagues. "The findings provide further evidence that high levels of [exercise](#) from a young age may be beneficial to individuals with CP," the researchers write.

## New Evidence that Exercise Helps Physiology and Functioning in CP

Dr. Runciman and colleagues performed detailed assessments of body composition in six elite-level Paralympic athletes with hemiplegic CP. Patients with this condition have neuromuscular impairment on one side of the body (left or right), caused by brain injury at a very young age.

The six study subjects, average age 23 years, were all experienced track and field athletes, training for six to 20 hours per week. They competed

at an elite Paralympic level, with 100-meter sprint times between 11.5 and 12.6 seconds.

A technique called dual-energy x-ray absorptiometry was used to perform measures of body composition, including [bone mineral density](#) (BMD) and [fat mass](#). The researchers also measured fat-free soft tissue mass (FFSTM)—an indicator of muscle and other lean body content.

The results showed similar values for BMD and fat mass on the affected versus unaffected sides of athletes with hemiplegic CP. That was in contrast to previous studies in sedentary young people with CP, who have reduced BMD on the impaired side.

The Paralympic athletes did have reduced FFSTM on the affected side. Compared to the unaffected side, FFSTM was about 15 percent lower in both the upper and lower limbs.

Overall, BMD levels for the Paralympic athletes were similar to those of age-matched able-bodied people. In fact, BMD values were higher for the athletes with CP compared to the general population, although the differences were not statistically significant.

Previous studies have reported reduced BMD and FFSTM in individuals with CP. "It is widely accepted that various physiological systems are impaired in individuals with CP as the result of brain damage at a young age and widespread neuromuscular dysfunction," according to the authors.

Adding to previous evidence, the new results suggest that in elite athletes with CP, the physiological processes affecting [body composition](#) are similar to those in able-bodied people of similar age. "These findings may highlight the importance of high levels of exercise training for all individuals with CP," Dr. Runciman and coauthors write.

In particular, the BMD findings in Paralympic athletes suggest increased density of cortical bone—the thick outer layer of bone, which responds to weight-bearing exercise. Training is also associated with normal fat mass, although the difference in FFSTM suggests a "ceiling" in how well the affected muscles can adapt to exercise.

"The combination of these findings have important implications for the involvement of children with CP in exercise as a management or rehabilitative tool," the researchers add. Studies have already shown that exercise improves functional performance and increases BMD in children with CP. The new findings in highly trained young adults with CP help in understanding how bone and other physiological systems adapt to exercise training.

"Thus there may be a significant number of children who have the capacity to participate in high-level [exercise training](#) from a young age," Dr. Runciman and colleagues conclude. However, they emphasize the need for follow-up studies to confirm the long-term benefits of weight-bearing exercise for young patients with CP.

**More information:** "Site-Specific Bone Mineral Density Is Unaltered Despite Differences in Fat-Free Soft Tissue Mass Between Affected and Nonaffected Sides in Hemiplegic Paralympic Athletes with Cerebral Palsy" [DOI: 10.1097/PHM.0000000000000532](https://doi.org/10.1097/PHM.0000000000000532)

Provided by Wolters Kluwer Health

Citation: Study of elite paralympic athletes supports benefits of exercise for children with cerebral palsy (2016, May 6) retrieved 1 May 2024 from <https://medicalxpress.com/news/2016-05-elite-paralympic-athletes-benefits-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.