

# Prevention better than fecal transplants

May 10 2016, by Cris Burne, Sciencenetwork Wa

---



Credit: CC0 Public Domain

Most people would think nothing of sharing a bed, knife or fork with their spouse, but would you ever share poo?

As gross as it sounds, faecal transplants are 90 per cent effective at treating recurrent gut infections, but UWA Professor Tom Riley says

preventing infection is poo-loads better than the cure.

Prof Riley has studied the potentially deadly *Clostridium difficile* gut infection ([CDI](#)) for 40 years, and says taking [antibiotics](#) to treat other infections is the biggest risk factor in inadvertently contracting *C. difficile*.

"Antibiotics kill all the normal good bugs in the gut, so *C. difficile* can get in there and do its thing," he says.

*C. difficile*'s 'thing' is causing infectious diarrhoea: in 2004 in Quebec, Canada, CDI caused more than 7000 CDI cases and 1270 deaths.

For decades, CDI has infected predominantly older, hospitalised patients, but Prof Riley says [virulent strains](#) are now infecting younger people in the wider community.

"In Australia we're in the midst of an epidemic of community-acquired infection," he says.

"These people don't get a severe disease, because they're inherently healthy, but my big concern is that these virulent strains will get into the hospitals, and then people will die."

## **Top-down or bottom-up?**

In adult humans, *C. difficile* grows best in a gut cleansed of its natural flora by antibiotics, so faecal transplants work by re-introducing billions of competing bacteria from a healthy gut.

"Generally speaking, you choose your spouse's poo, because your spouse's flora is most like yours," Prof Riley says.

"What's happened in the [USA], and even here, is to set up poo banks, like blood banks, where the whole process of screening for viruses and infections is done and the specimens are stored, ready to be transplanted.

"You have two options, the top-down approach, or the bottom-up approach, so you either do it via endoscopy or colonoscopy," Prof Riley says.

## Avoiding CDI

To minimise your chances of CDI, your best bet is to maintain a healthy biome.

"Avoid antibiotics if they're not really necessary... you have to have that insult to your [gut](#) flora before you're susceptible to C. difficile [infection](#)," Prof Riley says.

If you have been taking antibiotics, Prof Riley says it's important to wash with soap-and-water, as alcohol disinfectants won't kill C. difficile spores.

"Anywhere there's animal manure, you're likely to find C. difficile, so you need to be very careful about washing your hands and vegetables," he says.

*This article first appeared on [ScienceNetwork Western Australia](#) a science news website based at Scitech.*

Provided by Science Network WA

Citation: Prevention better than fecal transplants (2016, May 10) retrieved 25 April 2024 from <https://medicalxpress.com/news/2016-05-fecal-transplants.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.