

Frequent religious service attendance linked with decreased mortality risk among women

May 16 2016



Credit: George Hodan/public domain

Women who attended religious services more than once per week were more than 30% less likely to die during a 16-year-follow-up than women who never attended, according to a study from Harvard T.H. Chan School of Public Health. Frequent attendees also had significantly lower risk both from cardiovascular- and cancer-related mortality.

The study will be published online May 16, 2016 in *JAMA Internal Medicine*.

"Our results suggest that there may be something important about religious service attendance beyond solitary spirituality," said Tyler VanderWeele, professor of epidemiology at Harvard Chan School and senior author of the study. "Part of the benefit seems to be that attending religious services increases social support, discourages smoking, decreases depression, and helps people develop a more optimistic or hopeful outlook on life."

Nearly 40% of Americans report attending religious services once per week or more. Previous studies have suggested a link between attendance and reduced [mortality](#) risk, but many were criticized for major limitations, including the possibility of "reverse causation"—that only those who are healthy can attend services, so that attendance isn't necessarily influencing health. The new study addressed these criticisms by using rigorous methodology that controlled for common causes of attendance and mortality, used a larger sample size, and had repeated measurements over time of both attendance and health.

The researchers looked at data from 1992-2012 from 74,534 [women](#) who participated in the Nurses' Health Study. The women answered questionnaires about their diet, lifestyle, and health every two years, and about their religious service attendance every four years. The researchers adjusted for a variety of factors, including diet, physical activity, alcohol consumption, smoking status, body mass index, social integration, depression, race and ethnicity.

Compared with women who never attended religious services, women who attended more than once per week had 33% lower mortality risk during the study period and lived an average of five months longer, the study found. Those who attended weekly had 26% lower risk and those

who attended less than once a week had 13% lower risk.

The study also found that women who attended [religious services](#) once per week or more had a decreased risk of both cardiovascular mortality (27%) and cancer mortality (21%).

One limitation of the study is that it consisted mainly of white Christians and therefore might not be generalizable to the general population, other countries, or areas with limited religious freedom. In addition, the study population included only U.S. nurses of a similar socioeconomic status, who tend to be fairly health conscious.

"In this issue of *JAMA Internal Medicine*, Li et al report a clear and moderately strong association between attendance at religious services and decreased mortality during a 16-year follow-up of a subgroup from the Nurses' Health Study. ... First, readers and investigators must, as do these authors, focus on the data, no more and no less, and not attempt to generalize beyond the evidence. ... So what can we learn from this study? In this well-designed secondary data analysis, attendance at religious services is clearly associated with lower risk of mortality. This finding should not be ignored but rather explored in more depth," writes Dan German Blazer, II, M.D., M.P.H., Ph.D., of Duke University Medical Center, Durham, N.C., in a related commentary.

More information: "Association of Religious Service Attendance with Mortality Among Women," Shanshan Li, Meir J. Stampfer, David R. Williams, Tyler J. VanderWeele, *JAMA Internal Medicine*, online May 16, 2016, [DOI: 10.1001/jamainternmed.2016.1615](https://doi.org/10.1001/jamainternmed.2016.1615)

Commentary: *JAMA Intern Med*. Published online May 16, 2016.
doi:10.1001/jamainternmed.2016.1626

Provided by Harvard T.H. Chan School of Public Health

Citation: Frequent religious service attendance linked with decreased mortality risk among women (2016, May 16) retrieved 7 May 2024 from <https://medicalxpress.com/news/2016-05-frequent-religious-linked-decreased-mortality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.