


Middle East Respiratory Syndrome associated with higher mortality, more severe illness

May 18 2016



MERS is associated with higher mortality, more severe illness. Credit: ATS

Patients with Middle East Respiratory Syndrome (MERS) develop more severe critical illness and have higher mortality than patients with non-MERS severe acute respiratory infection (SARI), according to investigators involved with the largest study of critically ill patients with MERS. The study was presented at the ATS 2016 International Conference.

Until March 20, 2016, 1690 cases of MERS had been reported to the World Health Organization, with 80 percent of those cases in Saudi Arabia. The reported cases had an overall mortality of 35 percent. The investigators - led by Yaseen Arabi, MD, chairman, Intensive Care Department, and professor, College of Medicine, King Saud bin Abdulaziz University for Health Sciences and King Abdullah International Medical Research Center, Riyadh, Saudi Arabia - embarked on their research because few studies have examined the clinical course of critically ill [patients](#).

They collected data from 14 hospitals in 4 cities in Saudi Arabia and compared [critically ill patients](#) with laboratory-confirmed MERS to those admitted with SARI of non-MERS etiology during a similar period.

A total of 299 MERS patients with SARI were admitted to the participating ICUs and were compared with 218 patients with SARI only. Patients with MERS were younger (median age, 58 years) compared with non-MERS patients ([median age](#), 70 years). There were a number of chronic morbidities common in both groups, including

diabetes and liver disease. Symptoms such as cough, shortness of breath, and sputum production were also similar.

"Patients with MERS were more likely to be hypoxemic and to require invasive mechanical ventilation (85 percent versus 73 percent, $p = 0.001$), vasopressor therapy (77 percent versus 55 percent, $p = 0.001$), and renal replacement therapy (47 percent versus 23 percent, $p = 0.001$)," the investigators wrote. Mortality was higher in MERS versus non-MERS patients (70 percent versus 36 percent, $p = 0.001$).

"There is a substantial overlap in clinical presentation and co-mortalities among patients with MERS and SARI of other etiologies, making

More information: Abstract 9954: Critically Ill Patients With the Middle East Respiratory Syndrome (MERS): A Multicenter Study

Provided by American Thoracic Society

Citation: Middle East Respiratory Syndrome associated with higher mortality, more severe illness (2016, May 18) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-05-middle-east-respiratory-syndrome-higher.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--