

2016: Optimal imaging matches intravascular ultrasound for guiding PCI EuroPCR

May 17 2016

Using optimal frequency domain imaging (OFDI) to guide percutaneous coronary intervention (PCI) with second-generation drug eluting stents (DES) achieves equivalent clinical and angiographic outcomes to intravascular ultrasound (IVUS)-guided PCI at 12 months, show results from the OPINION study reported in a late-breaker session at EuroPCR 2016.

IVUS is an imaging modality used to assess lesion morphology during PCI and several studies have reported that IVUS guidance in PCI might be helpful to improve outcomes. OFDI is the most recently developed [optical coherence tomography](#) (OCT) technology that uses light rather than ultrasound to give high-resolution imaging.

"OPINION is the first study to compare clinical outcomes with OFDI and IVUS to assess lesion morphology during PCI with second-generation drug-eluting stents," explains study author Takashi Kubo, from Wakayama University, Wakayama, Japan.

The OPINION study researchers randomised 800 patients (mean age 68 years) scheduled for PCI using DES to either OFDI-guided PCI or IVUS-guided PCI at 42 centres in Japan. Patients underwent PCI using drug-eluting stents with resorbable polymer in de novo native coronary artery lesions. They were followed-up with angiography at eight months and the primary endpoint was target vessel failure assessing during clinical

follow-up at 12 months.

Results of the non-inferiority study reported at EuroPCR 2016 showed a similarly low rate of target vessel revascularisation, a composite of cardiac death, target vessel-related MI and clinically driven target vessel revascularisation, at 12 months in patients undergoing OFDI-guided PCI (5.2%) as in those randomised to IVUS-guided PCI (4.9%).

"The clinical outcome in both OCT-guided PCI and IVUS-guided PCI was excellent," reports Dr Kubo. He suggests the study results contribute to defining the clinical value of OCT guidance in PCI. "Our results might influence the next ESC guidelines, which currently give a class IIb recommendation for OCT. OCT use during PCI should have a class IIa recommendation," he says.

Commenting on the findings Takashi Akasaka, Professor of Medicine at Wakayama Medical University, Wakayama, Japan, says, "There has previously been no direct comparative data for OCT/OFDI-guided PCI and IVUS-guided PCI. The OPINION study succeeded in demonstrating that OCT-guided PCI resulted in equivalent clinical and angiographic outcomes at 12 months to IVUS-guided PCI."

Professor Akasaka considers the clinical outcome with both imaging strategies was very good, with a low rate of target vessel revascularisation. "Thus, the OPINION study may contribute to define the clinical value of the OCT guidance in PCI." He agrees that the findings mean that OCT/OFDI-guided PCI should be given a class IIa recommendation.

Provided by PCR

Citation: 2016: Optimal imaging matches intravascular ultrasound for guiding PCI EuroPCR

(2016, May 17) retrieved 3 May 2024 from <https://medicalxpress.com/news/2016-05-optimal-imaging-intravascular-ultrasound-pci.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.