

Study reveals safety and feasibility of robotically assisted PCI in complex cases

May 4 2016

A first-of-its kind study using robotic technology to remotely control coronary guidewires and stents reported on the feasibility of performing percutaneous coronary intervention (PCI) on patients with complex coronary lesions. Similar clinical outcomes compared to the PCI procedure performed manually were reported. Results of the 18-month complex robotically assisted percutaneous coronary intervention (CORA-PCI) trial were presented today as a late-breaking clinical trial at the Society for Cardiovascular Angiography and Interventions (SCAI) 2016 Scientific Sessions in Orlando, Fla.

The technology of performing robotically assisted PCI is relatively new and was recently approved by the FDA. Previous studies were performed largely on patients with simple coronary lesions. "For the first time, we've shown nearly identical outcomes with a group of very complex patients," said Ehtisham Mahmud, MD, FACC, FSCAI, the study's lead author and division chief, cardiovascular medicine and director, interventional cardiology and cardiac cath lab at the University of California, San Diego School of Medicine. He noted that approximately 80 percent of lesions treated in the CORA-PCI trial were complex, which is more reflective of contemporary interventional cardiology practice.

Consecutive robotic and manual procedures were performed by a single operator and a total of 334 PCI procedures were analyzed (108 R, 157 lesions; 226 M, 336 lesions) in 315 patients (age 67.7 ± 11.8). The majority of patients (78 percent) were male. Clinical success was similar

between the groups (99.1 percent for robotic PCI vs. 99.6 percent for manual PCI, $p=0.64$).

In the robotically assisted PCI group, over 90 percent of the procedures were performed entirely robotically or with minimal manual assistance. Procedure time was longer (robotic PCI $44:30 \pm 26:04$ min vs. manual PCI $36:34 \pm 23:03$ min, $p=0.005$) and remained significant after accounting for Syntax score, primary lesion length, and complexity ($p=0.026$); however, the longer procedure time was limited to the low complexity lesions. Propensity matched analysis ($n=94$ each group) confirmed the longer procedure time in the R-PCI group with comparable clinical success (98.9 percent vs. 100 percent, $p=1.0$). No increase in resource utilization or patient radiation exposure was reported with robotic PCI.

Furthermore, according to Dr. Mahmud, the data presented show that using robotic technology to perform PCI is a viable alternative to manual PCI with the potential benefits of reducing radiation exposure and orthopedic injuries to the interventional cardiologist.

Dr. Mahmud reports the following disclosures: Corindus, consulting and clinical trial support. No support received for this study.

Dr. Mahmud presented "Demonstration of the Safety and Feasibility of Robotically Assisted Percutaneous Coronary Intervention in Complex Coronary Lesions: Results of the Complex Robotically Assisted Percutaneous Coronary Intervention (CORA-PCI) Trial" on Wednesday May 4, 2016 at 1:30 p.m. ET.

Provided by Society for Cardiovascular Angiography and Interventions

Citation: Study reveals safety and feasibility of robotically assisted PCI in complex cases (2016,

May 4) retrieved 24 April 2024 from <https://medicalxpress.com/news/2016-05-reveals-safety-feasibility-robotically-pci.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.