

Survey: Common asthma warnings overlooked

May 11 2016



Dr. David Beuther examines an adult patient for breathing problems at National Jewish Health in Denver. One in 200 adults are newly diagnosed with asthma each year. Credit: National Jewish Health

A new national asthma survey commissioned by National Jewish Health shows that many adults are unaware of common symptoms of asthma in adults. Doctors say that the findings explain why many adults with asthma may not realize that they have the disease, and don't seek treatment that can help them.

The survey shows that, while people are aware of some of the most common [asthma](#) signs such as wheezing and shortness of breath, many don't recognize other warning [symptoms](#) like [trouble sleeping](#), [chest pain](#), and [persistent cough](#).

"A lot of people have asthma and don't know it. Many [adults](#) do not have the traditional asthma symptoms, or they don't have all of the symptoms," said David Beuther, MD, pulmonologist at National Jewish Health. "It's not rare that your asthma doesn't present like the textbook. It's actually more common than most people realize. To the patient or perhaps the primary care provider, things that seem like a very unusual set of symptoms for asthma are actually quite common. That is why you often need a specialist to diagnose and treat it."

Dr. Beuther said that one in every 200 adults are newly diagnosed with asthma every year. And while children with the disease might have more common symptoms, adults who develop asthma typically have some of the less recognized symptoms.

ASTHMA SURVEY

- Which are of the following are symptoms of adult onset asthma?
- Shortness of breath: 89%
- Wheezing: 85%
- Persistent cough: 65%
- Chest pain: 54%
- Trouble sleeping: 51%



Laurie Armijo of Niwot, CO uses a rescue inhaler after recently discovering that her persistent cough was actually adult-onset asthma. Credit: National Jewish Health

The survey, commissioned by National Jewish Health, was conducted by ORC International Telephone CARAVAN on March 3-6, 2016 via telephone. The combined sample consists of 1,002 adults (ages 18 years and older) living in the continental United States.

"These results are not surprising to those of us who see patients regularly. As a pulmonologist, I see people with symptoms that aren't yet defined like difficulty breathing, cough and episodes of recurring bronchitis. When we first embark on trying to figure out what that is, I often mention that one of the big three causes of chronic cough, for example, is asthma. And it's more common than not that my patient is surprised to hear that a chronic cough or recurring bronchitis is actually asthma," said Dr. Beuther.

An avid hiker and cyclist diagnosed with asthma in her forties, 54-year-old Laurie Armijo of Niwot, Colorado is one of the more than 17 million adults who have asthma.

"When I was diagnosed with exercise induced asthma it was very difficult because I had to pretty much give up a lot of the sports that I love. It was kind of scary for me because I could not breathe to run, walk fast, swim or ride my bike. I basically couldn't breathe," Armijo said.

Armijo now uses an inhaler when exercising so that she could resume her active lifestyle. Good medical care and monitoring can keep asthma under control.

Provided by National Jewish Health

Citation: Survey: Common asthma warnings overlooked (2016, May 11) retrieved 23 April 2024 from <https://medicalxpress.com/news/2016-05-survey-common-asthma-overlooked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.